

Dr. John C. Bugher, M.D., Director  
Division of Biology and Medicine

May 13, 1954

Gordon M. Dunning, Health Physicist  
Biophysics Branch, Division of Biology and Medicine

26383

ESTIMATED RADIATION DOSE TO THYROID OF NATIVES FROM RONGELAP

This memorandum is in reply to your request for an estimate of additional doses to the thyroid of the Rongelap natives due to the fact that tellurium, as a precursor to iodine, may be present in the gut after ingestion of fallout material. The tellurium, in turn, might disintegrate into radioactive iodine while in the gut, with subsequent deposition of the iodine in the thyroid.

There are some 17 radioactive isotopes of tellurium but only 7 of these are produced in fission. Of these, 6 are not of interest (4 have too short a half-life, 1 leads to stable iodine-127 and 1 leads to iodine-129 with a half-life of  $2.4 \times 10^7$  years). The remaining radioisotope is tellurium-132, with a half-life of 77 hours leading to iodine-132 with a half-life of 2.4 hours. (Incidentally there is no tellurium precursor that is of interest here.)

Without having the original data of LASL I have accepted their estimate that there were ingested and/or inhaled the products of  $5 \times 10^{13}$  fissions, assumed they were all ingested, and then proceeded to calculate the dose to the thyroid from (a)  $I^{131}$  (b) each short-lived iodine isotope of interest and (c) the added dose coming from  $T^{132}-I^{132}$ . The calculations show that  $T^{132}-I^{132}$  will produce an added dosage of about 26%.

The best estimated percentage absorption and deposition of iodine is yet to be determined. The best estimate I can turn up to date is still the 20% quoted in NBS Handbook 52. However, I will continue to search for additional information. In the meantime the table below indicates the magnitude of doses to the thyroid if one assumes 20, 50, and 100% absorption and deposition. Incidentally, it may be noted that the calculations below based on 20% (the number assumed by LASL) show estimated doses to the thyroid from  $I^{131}$  and from shorter lived iodine isotopes to be in good agreement with those estimated by LASL, i.e., 50 reps for  $I^{131}$  and 80 reps for short-lived iodine isotopes.

~~Information affecting the  
the United States within the  
Title 18, U.S.C.,  
any information or revelation  
obtained by any unauthorized person~~

US DOE ARCHIVES  
326 US ATOMIC ENERGY  
COMMISSION  
Collection ODM Files  
Box 3  
MRA T-S Radiation Exposures -  
Native and Test Personnel  
(March-May 1954)

CLASSIFIED, UN-CANCELLED  
BY AUTHORITY OF E.O. 12812, III-12 (ADMIN)  
BY *See Gray* DATE *3/26/81*  
BY *William A. Spencer* DATE *3-26-81*

OFFICE	BMBP				MILITARY RESEARCH & APPL. 7-5
SURNAME	DUNNING:MKack	<i>GMS</i>			
DATE	5-13-54				

DOSE TO THYROID (REPS)

	<u>Assuming 100% Retention</u>		<u>Assuming 50% Retention</u>		<u>Assuming 20% Retention</u>	
I <sup>131</sup>	255	255	128	128	51	51
I <sup>132</sup>	27*	(54)**	14*	(27)**	5*	(10)**
I <sup>133</sup>	370	370	185	185	74	74
I <sup>135</sup>	60	60	30	30	12	12
I <sup>132</sup> (Te <sup>132</sup> )	185*	(370)**	93*	(185)**	37*	(74)**
Total	897*	(1109)**	450*	(555)**	189*	(221)**

\* If assume that one-half of the I<sup>132</sup> (half-life 2.4 hours) present in the gut is deposited in the thyroid.

\*\* If assume all of the I<sup>132</sup> (half-life 2.4 hours) present in the gut is deposited in thyroid.

Most probable estimate of ratio of doses to thyroid is:

$$\frac{I^{132,133,135} + I^{132}(Te^{132})}{I^{131}} \approx 2.5$$

This material contains information affecting the national defense of the United States within the meaning of the Espionage Laws, Title 18, U.S.C., and the transmission or revelation of its contents in any manner to an unauthorized person is prohibited by law.

OFFICE ▶					
SURNAME ▶					
DATE ▶					

ANNEX

Calculations of Dose to Thyroid

<sup>131</sup>I

Assume: inhalation and/or ingestion of  $5 \times 10^{13}$  fissions

At D / 1 there are 0.017 d/e/10<sup>100</sup> fissions

or  $8.5 \times 10^7$  d/m/5  $\times 10^{13}$  fissions

or 38.3  $\mu$ c intake of <sup>131</sup>I

or  $1.4 \times 10^{12}$  atoms intake of <sup>131</sup>I

(Average energy 0.22 Mev)

Dose (reps) =  $\frac{(1.35 \times 10^{12})^*(0.22)(1.6 \times 10^{-6})}{(20)(93)} = 255$  reps

\* Correction for biological decay.

<sup>132</sup>I

Assume: inhalation and/or ingestion of  $5 \times 10^{13}$  fissions

At D / 1, <sup>132</sup>I intake is  $1.1 \times 10^{11}$  atoms

The average mean energy of <sup>132</sup>I is about 0.55 mev or 2.5 times that of <sup>131</sup>I.

Thus, the energy equivalent to <sup>131</sup>I would be

$(1.1 \times 10^{11})(2.5) = 2.75 \times 10^{11}$  atoms of <sup>131</sup>I

However, due to the short half-life of <sup>132</sup>I (2.4 hrs) assume that the energy equivalent of  $1.5 \times 10^{11}$  atoms of <sup>131</sup>I reaches the thyroid.

Thus, the ratio of doses  $\frac{I^{131}}{I^{132}} = 9.0$

This material contains information affecting the national defense of the United States within the meaning of the Espionage Laws, Title 18, U.S.C., and the transmission or revelation of its contents in any manner to an unauthorized person is prohibited by law.

OFFICE				
SURNAME				
DATE				

<sup>133</sup>I

Assume: inhalation and/or ingestion of  $5 \times 10^{13}$  fissions

At D / 1, <sup>133</sup>I intake is  $1.24 \times 10^{12}$  atoms

The average mean energy of <sup>133</sup>I is about 0.36 or 163 times that of <sup>131</sup>I.

Thus, the ratio of doses  $\frac{I^{131}}{I^{133}} \approx 0.7$

<sup>135</sup>I

Assume: inhalation and/or ingestion of  $5 \times 10^{13}$  fissions

At D / 1, <sup>135</sup>I intake is  $2.36 \times 10^{11}$  atoms.

The average mean energy of <sup>135</sup>I is about 0.3 mev or 1.36 that of <sup>131</sup>I.

Thus, the energy equivalent to <sup>131</sup>I would be


$$(2.36 \times 10^{11})(1.36) = 3.2 \times 10^{11} \text{ atoms of } I^{131} \text{ energy equivalent.}$$

Thus, ratio of doses  $\frac{I^{131}}{I^{135}} \approx 4.2$

~~CONFIDENTIAL~~

OFFICE ▶					
SURNAME ▶					
DATE ▶					





Assume: inhalation and/or ingestion of  $5 \times 10^{13}$  fissions.

At  $D \neq 1$ ,  $\text{Te}^{132}$  intake 100  $\mu\text{c}$

or  $1.5 \times 10^{12}$  atoms

Assume: the time spent in the gut is 77 hrs.

Then,  $7.3 \times 10^{11}$  atoms of  $\text{Te}^{132}$  will have disintegrated into  $7.3 \times 10^{11}$  atoms of  $\text{I}^{132}$ .

The average mean energy of  $\text{I}^{132}$  is about 0.55 Mev or 2.5 times that of  $\text{I}^{131}$ .

Thus, the energy equivalent to  $\text{I}^{131}$  would be

$$(7.3 \times 10^{11})(2.5) = 1.8 \times 10^{12} \text{ atoms of } \text{I}^{131}.$$

However, due to the short half-life of  $\text{I}^{132}$  (2.4 hrs), assume that only the energy equivalent of  $1 \times 10^{12}$  atoms of  $\text{I}^{131}$  reaches the thyroid.

Thus, the ratio of doses  $\frac{\text{I}^{131}}{\text{I}^{132}(\text{Te}^{132})} \approx 1.4$ .

DISTRIBUTION: copy 1A - Dr. Bugher  
 2A - Dr. Fanning  
 3A - Biophysics Branch  
 4,5,6A - BOM Files

*Cy 5-A destroyed 6/14/54*

~~Contains information affecting the national defense of the United States within the meaning of the espionage laws, the transmission or the revelation of its contents in any manner to an unauthorized person is prohibited by law.~~

OFFICE ▶					
SURNAME ▶					
DATE ▶					