Itt I may

AFOAT-1/TD-3 370.009

SUBJECT:

Height of IV!

Lilie Cloud

TO:

Commanding General
Air Research and Development Command
ATTY: Col. R. M. Isbell
Fost Office Lox 1395
Ealtimore 3, Maryland

- l. Reference your communication of 27 February 1953, subject as above, and telephone communication between Col. Isbell and Dr. Urry of this office delaying reply until further discussion with Lt. Col. Lulejian, reply to your request is attached as Inclosure 1 hereto.
- 2. As this subject is somewhat outside of the scope of the mission of this office, the data transmitted and the views expressed by Dr. Urry are not necessarily official views of Headquarters USAF, AFCAT-1.

BY COLLIAND OF THE CHIEF OF STAFF:

Copy to Dr. Nm. Ogle, LASI.

HARVEY W. C. SHELTON Colonel, USAF Executive, AFOAT-1 Office for Atomic Energy DCS/0

RG 326 US ATOMIC ENERGY COMMISSION

Collection J DIU.

Folder OP. IVY GENERAL,
TEST PLANNING

BY R.G. BOGER DATE:

Copied/DOE



HEADQUARTERS USAF OFFICE FOR ATOMIC EMERGY, DCS/G AFOAT-1

2H# 17473

4 MAY 1953

24 April 1953

MELIORANDULI FOR RECORD

SUBJECT:

Height of IVY (Recommended)

Mke Cloud

- 1. At the time of the IVY kike Operation two aircraft designated "Saltshaker" flew race-track courses, one due south of ground zero at approximately 70 nautical miles, the other due east of ground zero at approximately 80 nautical miles. The planned mission of these flights was to secure a photograph of the IVY Mike cloud each minute for one hour following the explosion. This mission was requested by Headquarters USAF, AFOAT-1 for after-the-fact cloud height calculations but the pictures have not yet been received, and may not be of much value for the intended purpose because of pocrly defined or absent horizons.
- Unscheduled bubble sextant readings were made by Dr. W. D. Urry in the aircraft to the south and Colonel Fee of Headquarters USAF, AFCAT-1 in the aircraft to the east. Aircraft loran positions were provided by the navigator of the aircraft who also checked some of the angle observations. Attached hereto is Table I giving the observations and calculations of cloud height and heights of various outstanding features of the cloud. The observations of angle, distance to ground zero, and aircraft altitude are believed to be such that the calculated heights are not in error one way or the other by more than a few thousand feet. Consideration of some points would lead one to believe that no large errors were introduced by "edge" sighting and thereby obtaining erroneously high angles. First of all, there can be little of this kind of error in sighting on the rather sharp-pointed plume which yielded an altitude of 135,500 feet. The difference between this altitude and the top of the cloud at around 120,000 feet was in the correct proportion to the thickness of the cloud (120,000 - 67,000 = 43,000 feet) as judged at the time. Secondly, a sighting on the far right edge of the cloud (cbs: at 11.75 minutes in Table I) gave 104.000 feet and this can hardly be in error by 40.000 feet due to erroneous sighting arising from edge and thickness effects, as it would be if the cloud did not rise above the tropopause. Thirdly, the main shear layer measured at 15.25 minutes in Table I gave an altitude corresponding exactly to that of the prognosticated principle shear altitude for the event and was again judged to be at the correct propertional altitude for a top at around 120,000 feet. Fourthly, the altitudes of 110,000 and 112,000 feet at 2.66 and 3.42 minutes are not explainable on false base line because of lateral movement of the cloud; air movements are not that rapid compared with a plane to ground zero base line of 68 nautical miles.

Inclosure 1 1

Copied/DOE


- 3. The constancy of the measured height of the top of the cloud over a period of one hour in Table I is consistent with the slow movement of the cloud. Measurements by Colonel Fee in the aircraft to the east of ground zero are not so consistent and calculated heights appear to decrease with time, probably due to an increase in true base line owing to a west-ward movement of the cloud. Colonel Fee's data place the top of the cloud variously between 127,000 and 99,000 feet.
- It is to be noted that a marked vertical deceleration occurred at two minutes, at which time the conspicuous lateral spreading occurred. This is in agreement with the observations of C. E. Falmer except that actual measurement indicates the "splashing" occurred against a barrier between 100,000 and 110,000 feet, which may be of significance in connection with the usual increase in temperature at around 100,000 feet. Some observations given to Dr. Urry by Col. Morris in the B/36 sampling control aircraft are plotted also in Figure I. Col. Morris's observation at 40 seconds as the cloud went by his altitude is in fair agreement with Dr. Urry's curve as also is his observation some time after 20 minutes. At five minutes, Col. Morris's observation is considerably lower than Dr. Urry's but the top is well within the stratosphere.
- 5. Some observations by Lr. Singlevich of Headquarters USAF, AFCAT-1 under similar conditions, but at 30 to 40 nautical miles from ground zero at the time of IVY King test are shown on Figure I for comparison.
- 6. The hypothesis put forward by Professor C. E. Palmer in his second letter of 2 February 1953 for the secondary formation of the main large cloud appears to the author to be plausible but it also appears equally certain that the bulk of this main cloud ended up in the stratosphere between 60 and 120 thousand feet. The maintenance of its shape and form are believed to be due to its internal turbulence and unequal heat distribution. Any estimate of the degree of mixing of the secondarily formed large main cloud with a primary column of nuclear debris, if indeed this did not intimately occur during rise, is purely hazardous guesswork. As far as is known, no observations were made for guidance on this aspect of the problem. It seems apparent that sampling was conducted some thousands of feet below the bottom of the main cloud, as statements by some of the pilots of the F-E4 sampling aircraft, appears to confirm.
- 7. There is attached hereto as an integral part of this problem a memorandum from Mr. P. W. Allen to Dr. W. D. Urry dated 11 December 1952, discussing a well-defined relation between early rate of rise of the cloud and the energy of a nuclear explosion.



ti. D. URMY
Assistant for Nuclear Physics
Office of the Technical Director, AFCAT-1

3 Incls
Table 1
Figure I
Vemo dtd 11 Dec 52 fm Pr. Allen to Dr. Urry

Copled/DOE LANL, J-DIV.



100



CLOUD HEIGHT DATA FOR WIKE SHOT, OF STION IVY, Taken by Dr. W. D. Uhry

In Local Hov 52	Decimal Time M / Min	Aircraft Position Latitude Longitude	Aircraft Altitude Fest	Distance i Ground Naut. Mi.		Elevation Angle Deg Min	Calculated Height Above Aircraft	Correction for Earth Curvature	Total Height	Coordinates of Ground Zero Approx. 11-40.2 N 162-11.7 E
F3-4		10-36 N							•	
0715	0	161-17 E	12300	68,6	416000					
0716.5	1.5	10-35 N 161-50 E	12300	68.5	415000	05-47	42200	4000	58200	Absolute Too, Mair
	Anderson State of Sta	10-35 M								~ ານເ
0717.7	2.66	161-52 E	12300	67.9	412000	12-50	94000	4000	110000	
0718.4	3.42	10-34.5N 161-53.5E	12300	67.9	412000	13-04	75300 ·	4000	111800	
		10-34.5N								
0719.4	4.42	161-55-5E 10-31-25M		67.6	411000	13-33	99000	4000	115000	Top of main cloud,
0720.8	5.75	161-57 E	12300	67.5	41,0000	14-10	103500	4000	119500	plume above top of
0723.5	8.5	10-37 H 161-50 E	12300	66.9	405500	16-28	119500	4000	135500	Point A, Picture / Top of plume
Y.C. RANGE		10-39.5N	LANGE VIN MANAGEMENT	ayan waxan a Kalifalla baran asaa asaa	A STATE OF THE PARTY OF THE PAR		and the state of t	Total Control of the	Andrew Property and the Party	Point B. Picture /
0725.5	10.5	161-45.25E	12300	65.9	400000	07-18	51200	4000	67200	Base of musbroom
0706 0	11 78	10-40.5H 161-44 E	12300	65.5	2075.00	12-30	82000	4000	104000	Point C, Picture L
0726.8	11_75	10-41.5N	16360	325	397500	12-30	60000	4000	104000	Right side of ton Mair
0728.4	13.4	161-41.75E	12300	55.8	399000	14-38	104000	1000	120000	Absolute top. Clou
0730.3	15.25	10-42.5H 161-39.5E	12300	65.9	400000	01-35	11100	1000	27100	Point D, Picti C Lower shear laver
MINAN		10-38.5N		<u>XaZ 0.7</u>	TANANA.		-			THE SHIPE AND A
0811	56	161-35 E	12300	71.1	-132000	14-00	109000	-4000	124000	Top of orange cle
0812	57	10-38.5N 161-33 E	12300	72.5	440000	13-55	109000	-1000	125000	Top of orange sla
EX.B.X.		10-33 N			ann agus An is Carles Mallace					
0815	60	161-24 E	12300	78.0	474000	_13-17	112000	4000	128000	Top of coares Car
0818	63	10-34 N 161-29-5E	12300	77.8	472000	16-35	140000	4.000	156000	But the same for a
	65	10-32.5N		78.1	475000	12-10-	102000	~ 1000	118000	***
0820		161-31.5E		Kthan	315000	44_4				- ;;
								RESP 30	DATE "	· · · · · · · · · · · · · · · · · · ·

This Document Consists of _



17.Allen/whk/62189

SUBJ: The Relationship Between Energy and Rate of Cloud Rise

TO: Dr. N. D. Urry

FRCE: P. W. Allen

DATE: 11 Dec 52

71.17

- l. As you suggested last week, I have gone over all available data on the rates of rise of clouds from U.S. atomic tests and have plotted values for the rates of rise against energy in EKT, as shown in the accompanying graph, together with the empirical equation of relationship.
- 2. The rate of rise changes with time, increasing to a maximum during the first minute and decreasing thereafter to essentially zero after about 10 minutes. The data available to us are not good enough to show the maximum rate due to poor timing and infrequent measurements, but may be used to obtain the average rate off rise over a period of minutes. The average over the initial 3 minutes is used on this graph. In all cases except IVY like the clouds were still rising rapidly and were still in the troposphere after the third minute. The like cloud was treated in a special manner as indicated below.

It is reasonable to believe there to be some dependency of rise rate on the lapse of ambient air temperature with altitude. The offect of inversions and stable layers will, however, be a minimum in the earliest seconds of rise, increasing in importance as the temperature difference decreases between cloud and surrounding atmosphere. When the cloud reaches ambient air temperature, further vertical motion is damped out. It is therefore preferable to measure the rate of rise at the earlies possible time, and the maximum rate of rise should be more indicative of energy than the mean 3-minute rate used here.

Since the mean lapse rate of temperature is markedly different in the stratosphere than in the troposphere, it is preferable to make all measurements in the troposphere until adequate corrections can be made for this.

- 3. Rate of rise data are available from the following sources:
- a. Overation SANDSTONE. Ir. Paul Humphreys, USNB, documented the rise and dispersion of the SANDSTONE clouds in an AFSNP publication, "Classified Scientific Meteorological Information, Operation SANDSTONE." His data were obtained by theodolite and are reasonably accurate over at least the first few minutes of rise
- b. Operation GPEENHOUSE. The rise of the GREENHOUSE clouds were obtained from an unpublished report on "Cloud Physics", Proj. 4.6, by Dr. W.W.Kellogg, Rand Corp. Motion picture photography were analyzed for cloud rise and cloud dimensions, and the rates of rise over the first 4 or 5 minutes are probably good, although weather clouds obscured parts of the atomic clouds. The maximum altitudes of the Dog and George clouds are still in doubt since the tops of these clouds were not visible from the camera positions.
- c. Operations BUSTER-JANGIE. Two sources of cloud rise data are available for these operations, one being that taken by myself (with your help in a

(5.

Incl 113

Copled/DOE





couple of cases) by hand clinometer, and the other taken by Air Weather Service personnel by theodolite. These two sets checked very well in all cases except Charlie cloud. On the graph, the average of the two is indicated by the circled dot, with the outriggers showing the values themselves.

d. Creration IVY. Three surface vessels and three aircraft were engaged in making cloud rise and height measurements of Like cloud. Of these, one vessel failed to make any height measurements until H + 2½ min., and one airplane mad only one measurement, at approximately H + 40 sec, before H + 5 min. It is not believed that timing was very accurate on this measurement so it was discarded. The other measurements showed the cloud to have approached maximum altitude at 3 minutes so in addition to the 3 minute average an average was obtained using earlier measurements. Since there is reason to believe the rate to decrease in the stratosphere (above 58000 ft on like day) the second average was taken of observations below that height. The two averages are shown as horizontal lines on the graph.

(1) Three minute heights:

U.S.S Curtiss	100,500 ft.
U.S.S.Rendova	117,000 ft.
Aircraft No. 1 (Dr. Urry)	111,000 ft.
Aircraft No. 2 (Col. Fee)	127.000 ft.
Average	114,000 ft. in 3 min.

(1) Extrapolation of troposphere rates:


U.S.S. Curtiss	46,600 ft. at 1 min.	139,800 ft/3
U.S.S. Rendova	He early measurements	
Aircraft No. 1	58,200 ft. at 12 min.	116,400 ft/3
Aircraft No. 2	No early measurements Average	128,100 ft/3

Three surface vessels and two aircraft made measurements of the King clou and all data are on hand except that from one surface vessel. The three minute heights are as follows:

U.S.S. Oak H111	58,300
U.S.S. Rendova	56 ,1 00
Aircraft No. 1 (Ltr. Singlevich) 58,300
Aircraft No. 2 (Col. Morris)	53,000 (Doubtful)
Average	56,400

The average is plotted on the graph.

4. Considerable improvement in this relationship might well result from more accurate determination of the rate of rise and in particular of the maximum rate, and from development of a correction factor for variations in the ambient air lapse rate of temperature.



P. W. ALLEN

