

400307

T A B

D

DOE ARCHIVES

5002485

CONCENTRATION OF <sup>137</sup>CS IN SUBSISTENCE CROPS AND FISH AT BIKINI ISLAND

FOOD PRODUCT	NO. OF SAMPLES	AVERAGE CONCENTRATION PCI/G NET WEIGHT	RANGE OF CONCENTRATION PCI/G NET WEIGHT
COCONUT MEAT (GREEN)	6	22.7	3.5-48
COCONUT MEAT (INTER-MEDIATE)	9	16.5	4.8-32
COCONUT MEAT (MATURE)	31	30.9	5.3-117
COCONUT MEAT (SPROUTED, SPRING)	8	27	16-52
ALL COCONUT MEAT	54	27	3.5-117
COCONUT FLUID	28	13.5	1.2-44
BREADFRUIT	2	6.5	5.2-7.8
SQUASH	12	8.5	1.6-20
PAPAYA	18	14	1.6-31
BANANA	3	0.92	0.54-1.3
SWEET POTATO	2	3.6	2.3-5
WATERMELON	17	2.6	0.26-7.2
GARDEN FRUITS AND VEGETABLES (AVERAGE OF SQUASH, PAPAYA, BANANA, SWEET POTATO, WATERMELON)		5.9	
FISH (MULLET) <sup>+</sup>	6	0.026 <sup>+</sup>	
DOMESTIC MEAT		15 <sup>*</sup>	

DOE ARCHIVES

+ FROM V. NOSHIKIN

ESTIMATED FROM BIKINI FIG DATA

5002486

CONCENTRATION OF <sup>90</sup>Sr IN SUBSISTENCE CROPS AND FISH AT EREU ISLAND

FOOD PRODUCT	NO. OF SAMPLES	AVERAGE CONCENTRATION PCI/G NET WEIGHT	RANGE OF CONCENTRATION PCI/G NET WEIGHT
COCONUT MEAT	9	0.021	0.0033 - 0.052
COCONUT FLUID*	-	0.021*	-
BREADFRUIT	2	1.9	0.47 - 3.4
WATERMELON	8	0.031	0.012 - 0.053
SQUASH	6	0.054	0.024 - 0.15
PAPAYA	5	0.29	0.052 - 0.39
SWEET POTATO	1	0.13	-
GARDEN FRUITS AND VEGETABLES (AVERAGE OF WATERMELON, SQUASH, PAPAYA, SWEET POTATO)		0.13	
FISH (MULLET)		0.076 <sup>+</sup>	
CLAMS		0.005 <sup>+</sup>	
DOMESTIC MEAT		0.011 <sup>**</sup>	

\* ASSUMED TO BE THE SAME AS COCONUT MEAT

+ FROM V. NELSON AND B. SCHELL

\*\* FROM 1975 BIKINI DOSE ASSESSMENT

DOE ARCHIVES

5002487

CONCENTRATION OF  $^{239+240}\text{Pu}$  IN SUBSISTENCE CROPS AND FISH AT ENEU ISLAND

FOOD PRODUCT	NO. OF SAMPLES	AVERAGE CONCENTRATION PCI/G NET WEIGHT	RANGE OF CONCENTRATION PCI/G NET WEIGHT
COCONUT MEAT	9	$2.8 \times 10^{-5}$	$4.1 \times 10^{-6}$ - $5.3 \times 10^{-5}$
COCONUT FLUID	-	$2.8 \times 10^{-5}$ *	-
BREADFRUIT	1	$1.7 \times 10^{-5}$	-
WATERMELON	8	$1.3 \times 10^{-5}$	$4.4 \times 10^{-6}$ - $2.0 \times 10^{-5}$
SQUASH	6	$8 \times 10^{-6}$	$3.5 \times 10^{-6}$ - $1.9 \times 10^{-5}$
PAPAYA	3	$8.3 \times 10^{-6}$	$6.5 \times 10^{-6}$ - $1.1 \times 10^{-5}$
GARDEN FRUITS AND VEGETABLE (AVERAGE OF WATERMELON, SQUASH, PAPAYA)		$9.8 \times 10^{-6}$	
FISH (MULLET) <sup>+</sup>	6	$1.3 \times 10^{-4}$ +	

\* ASSUMED TO BE THE SAME AS COCONUT MEAT

+ FROM V. NOSHKIN

DOE ARCHIVES

5002488