

COPY

404120

20833

DEPARTMENT OF DEFENSE
ARMED FORCES SPECIAL WEAPONS PROJECT
P.O. Box 2610
Washington, D. C.

June 25, 1952

Mr. Seth Woodruff
Las Vegas Operations Office
Mercury, Nevada

Dear Seth:

Conversations with Doctor John Bugher just before his departure for some well-earned leave indicated that the following dosages are acceptable for the roll-up operations phase:

a. Total of 3 roentgens, gamma only, for a ten-week period.

b. Ten milliroentgens non-contaminated area. Areas with instances of greater than 10 milliroentgens would be considered as contaminated, and a monitor should be present during operations in this area.

c. Two milliroentgens - level for vehicle and equipment contamination. This was the same level utilized during TUMBLER/SNAPPER.

These levels should allow reasonable roll-up operations without an undue expenditure of personnel. The important level is of course the total dosage of 3 roentgens for a ten-week period. This complies with accepted peacetime tolerances of .3 roentgen per week. Dr. Bugher promises to send this information over his signature when he returns from leave.

Sincerely,

/s/ Jim

James B. Hartgering
Lieutenant Colonel, MC
Assistant Surgeon

cc: Dr. Bugher
CO Test Cmd, AFSWP

COPY

DOE ARCHIVES