

STANDARD FORM NO. 64

Office Memorandum • UNITED STATES GOVERNMENT

TO : Dr. Bugher

DATE: 4/12/1953

FROM : F. Western

334

SUBJECT: Notes on Tripartite Conference

Attached are some notes on the Tripartite Conference as you requested. They are mostly from memory and may not be completely reliable. However, as a supplement to the notes and remember facts you have in hand, they may be of value. If there are particular points on which you wish confirmation, I will be glad to try to get it for you.

I am not certain of the nature of the report for which you plan to use the notes. Miss Laner has suggested that this is an appropriate topic for the six months report to the JCAE, for which copy is due April 17. Shall I write it up for that purpose or, since you are already engaged in reporting the meeting, would you prefer that your report be used also as a basis for a JCAE item?

fw

US DOE ARCHIVES	
326 U.S. ATOMIC ENERGY	
COMMISSION	
RG	<u>DOE HISTORIAN (DBM)</u>
Collection	<u>1132</u>
Box	<u>3363</u>
Folder	<u>#11</u>

Notes on Tripartite Conference, March 30-April 1.

The following notes are mostly from memory. In view of the fact that a stenographic record was taken, I did not anticipate the need of detailed notes. In some cases the statements may be unreliable.

March 30.

A.1. Some discussion of "rep" but no recommendations as to size or name.

2. There were some revisions in the "standard man" but I made no record of details.

3. Failla presented certain recommendations given in the draft of the report of the NCRP Sub-committee on External Dose. As I recall, the action was approximately as follows:

a. The recommendation that the permissible dose for persons over 45 years of age was discussed and strongly opposed by Morgan and, I believe, Marley. No action.

I believe that the recommended differential factor of 10 between children and adults was adopted.

b. I believe that Failla's recommendations for maximum permissible dose to local areas were accepted. These are:

For hands and forearms, feet and ankles, or head and neck, for any ionizing radiation, 1500 millirems to the skin provided that the dose to any other tissue shall not exceed that which would result in exposure of the skin to 1500 mr of ordinary x-rays, and provided that the exposure to the eyes shall not exceed 450 mr/week for ordinary x-rays nor 300 millirems per week for ionizing particles.

4. The currently accepted value of 10 for the r.b.e. of protons was reconfirmed for specific application to the lens of the eye (protection against radiation cataracts). Discussion indicated that the value could be much less for other applications but I recall no action on this point. The following maximum permissible values of incident neutron flux, interpolated from the results of the calculations of Mitchell, Snyder, and possibly others, based on an r.b.e. of 10, were adopted:

Energy (Mev)	10	5	2	1	0.5	0.1	0.01	Thermal
Flux (cm ⁻² sec ⁻¹)	30	30	40	60	80	200	1,000	2,000

These are average fluxes for a 40 hour week.

A uniform value of one for the r.b.e. of betas and recoil electrons was reconfirmed although it was noted that this probably represents an over simplification.

After apparent unanimous agreement that the r.b.e. of alpha particles for known effects in the body appears to be less than 5, insistence of some two or three persons for a more "conservative" value of 10 prevailed (?).

- A.5. I recall no definite action on genetic considerations.
6. I believe it was agreed to recommend limitation of dose to large populations by a factor of ten (?). This may be confused with B.6. March 31, ~~XXX~~ below, since I cannot recall two separate discussions on the subject.
7. No agreement was reached on length of time for averaging dose. The British were more conservative than the Americans, partly from administrative considerations and partly from the feel^{ing} by Mitchell that a system of averaging dose should include a six-months period of recovery.
- B. Except as included in the above notes, I recall no discussion on these items.

March 31.

- A. To conserve time, the first 5 items were passed over. However, item 5 came up for discussion later in connection with discussions of inhalation of uranium and thorium. It was agreed that the retention figure for insoluble dusts should be lowered from 25% to 10%. I do not recall the convention adopted for the disposition of the remaining 90%.
6. I believe it was agreed that a factor of 10 should be applied in reducing maximum permissible levels of internal dose to levels satisfactory for large populations. See A.6., March 30.
7. Possible effects of iodine on sheep were discussed and data presented by Healy. Marley was particularly concerned over the possibility of damage to local grazing animals. Dr. Warren presented data from effects of iodine on human euthyroids to indicate that the thinking of Hanford and the English on this subject is ultraconservative. Both seemed to feel that consideration of deposition on vegetation presented a more stringent limitation on air concentrations by a factor of more than 1,000 than does inhalation by humans.
- B. 1. Radium was discussed but no changes in current values were recommended. Levels of radon~~um~~ in air higher by a factor of 10 (?) than those currently accepted were recommended. I believe that these may be interpreted as radon in equilibrium with its decay products. (This will be a boon to the AEC and PHS in connection with the uranium mines.)
2. The maximum permissible body content for polonium was raised to 0.02 (?) uc. The British had been unable to reconcile data given by Flnk, NNES VI-3, with ~~XXXXXXXXXXXX~~ biochemical data used by Morgan in computing maximum permissible levels in air and water, and no action was taken on these (?).
3. ~~Actinium~~ Actinium, maximum body content, 0.02 uc. (?)
4. Thorium, maximum concentration in air same as for uranium.

- B.5. Plutonium. Studies of 15 cases of injections were reviewed by Langham. Urinary excretion was surprisingly uniform from one case to another and from one day to another. Excretion rates: at 1 day, 0.8%; 5 days, 0.3%; 9 mos, 0.003%, 1260 days, 0.001%, fitting equation

$$y_u = 0.23 x^{-0.77}. \text{ Versene increases rate by 10 to 50x.}$$

British discussed a case in which there is some evidence that as much as 50 mg was swallowed.

I recall no action on present values.

6. ??

7. The MPC of $C^{14}O_2$ in air was increased by a factor of 10 (?) to 1×10^{-5} uc/cc² on statement by Brues that less than 10% (?) of C^{14} in inhaled CO_2 enters into metabolism. There may have been other action on C^{14} which I do not recall.

8. The MPC of H^3HO in the atmosphere was reduced by a factor of 2 on the basis of Pinson's observation that the body absorbs as much through the skin as through the lungs.

April 1.

C. Failla's recommendations on accidental exposure were discussed extensively. The British were strongly opposed to his concept of radiation status unchanged by accidental exposures over maximum permissible levels. No agreement.

The British position was based largely on administrative considerations.

→ It was agreed that for bone-seeking radioisotopes, Calcium and Barium would be compared with Strontium to permit an average weekly dose to the skeleton of $0.3/5$ rps, while ~~not~~ others would carry an additional safety factor of 2.5 on account of possible differences in distribution to permit an average dose to the skeleton of $0.3/12.5$ rps per week. Hamilton aided the British in putting over the additional 2.5 which was opposed by Morgan, Marinellio ~~others~~