

410315

Reprinted from "Biological Aspects  
of Radiation Protection"  
edited by T. Sugahara and O. Hug,  
IGAKU SHOIN LTD., Tokyo, 1971.

## HEMATOLOGICAL EFFECTS ON HEAVILY IRRADIATED JAPANESE FISHERMEN

TOSHIYUKI KUMATORI

*Division of Radiation Health,  
National Institute  
of Radiological Sciences  
Chiba, Japan*

BEST COPY AVAILABLE

1971



IGAKU SHOIN LTD. TOKYO

## HEMATOLOGICAL EFFECTS ON HEAVILY IRRADIATED JAPANESE FISHERMEN

TOSHIYUKI KUMATORI

*Division of Radiation Health,  
National Institute  
of Radiological Sciences  
Chiba, Japan*

### INTRODUCTION

On 1 March in 1954, a thermonuclear test explosion was performed by the United States Authorities at Bikini Lagoon in the Pacific Ocean. At that time one Japanese fishing boat "the 5th Fukuryu-maru" was located at about 90 miles east from explosion center. The boat was nearly 100 tons and 23 fishermen aged from 18 to 39 were on board.

At about 3:40 a. m., while they were occupied in fishing tuna they saw a huge red light in the west and heard a detonation several minutes later. At about 7:00 a. m. white material began to fall, which continued for approximately 5 hours. They gave up fishing and returned to their mother port on 14 March 1954.

After landing, all the fishermen were found to have been injured by the radioactive materials. Seven of them were hospitalized to the Tokyo University Hospital and the other 16 were received by the First National Hospital of Tokyo by 28 March. They were discharged from both hospitals in May 1955, except for one fatal case who died on 23 September 1954. After being discharged we continued the follow-up studies so far as possible on annual basis. However, mainly because of the varied status of their occupations and widely distributed addresses, it was not always possible to get them all together for medical examination.

Medical data of the fishermen have been reported several times (MIYOSHI and KUMATORI, 1955; KOYAMA et al., 1955; MIKAMO et al., 1956; MIYOSHI and KUMATORI, 1962; KUMATORI and MIYOSHI, 1963; MIYOSHI and KUMATORI, 1964; KUMATORI et al., 1965).

### FALLING OF RADIOACTIVE MATERIALS

During the falling of the white materials, when its intensity was greatest, these persons were unable to keep their mouths and eyes open. Their footprints were clearly marked on the deck covered by fallout.

The fishermen were irradiated from three sources:  
(1) From the radioactive fallout on the deck etc.  
(2) Externally from the fallout on the deck etc.  
(3) Internally from the fallout on the deck etc.

The diameters of the fallout particles ranged from 0.5 to 10  $\mu$  (Kumatori et al., 1954). According to the measurements on 1 March, rare-earth elements activity and uranium content were determined. The specific activity of the material was estimated from these data, a value of 1.5  $\mu$  Ci/g. was obtained at 1 a. m. on 1 March. The estimated internal dose from the internally deposited radioisotopes was 1.5 r.

However, radiation from the fallout caused skin lesions. External radiation dose for 14 days was estimated to be 1.5 r. The percentage of these doses was reported on the results of the experimental and theoretical decay curves of many pl

Table 1. Estimated

Subject No.
T-1
T-2
T-3
T-5
T-6
T-7
T-8
K-1
K-2
K-3
K-4
K-5
K-6
K-7
K-8
K-9
K-10
K-11
K-12
K-13
K-14
K-15
K-16

\* T-3 put the fall-out amount added in total.

## N HEAVILY FISHERMEN

### STATE OF IRRADIATION AND ESTIMATED RADIATION DOSE

The fishermen were irradiated in the following 3 ways:

- (1) From the radioactive materials adhering to the body surface.
- (2) Externally from the radioactive materials deposited in the cabins, on the deck etc.
- (3) Internally from the radioactive materials entering various organs.

The diameters of the fallout materials, which were the powders of coral-reef fragments, ranged from 19 to 460  $\mu$ , mainly 100 to 400  $\mu$  (KIKUCHI et al., 1954). According to the results of radiochemical analysis of fallout on 26 March, rare-earth elements contributed about 50 per cent of the total radioactivity and uranium contributed about 20 per cent (KIMURA et al., 1956). The specific activity of the material was 0.37 mCi/g on 23 April. By extrapolation of these data, a value of 1.4 Ci/g was obtained as the specific activity at 7:00 a.m. on 1 March. The estimation of radiation dose, especially that from internally deposited radioactive materials was difficult.

However, radiation from body surface was large enough to cause skin lesions. External radiation dose was estimated as shown in Table I. The exposure dose for 14 days ranged from about 170 to 690 R, and nearly 60 per cent of these doses was received on the first day. This estimation was based on the results of the experimental reproduction of ash-fall, extrapolation of decay curves of many places of the boat, and on detailed investigation of

Table 1 Estimated dose of whole body gamma radiation.

Subject No.	First Day	Total
T-1	240~290 R	450~500 R
T-2	210~260	390~440
T-3	150~200	260~310* (360~410)
T-5	400~430	660~690
T-6	130~180	200~250
T-7	140~190	220~270
T-8	310~360	520~570
K-1	190~220	310~340
K-2	130~180	200~250
K-3	140~190	230~280
K-4	120~170	190~240
K-5	140~190	220~270
K-6	180~230	300~350
K-7	230~280	340~390
K-8	220~270	380~430
K-9	310~360	550~600
K-10	140~190	230~280
K-11	120~170	170~220
K-12	100~150	170~220
K-13	250~300	370~420
K-14	420~500	510~590
K-15	140~190	210~260
K-16	120~170	190~240

\* T-3 put the fall-out material close to his bed. Therefore about 100 R should be added in total.

osion was performed by  
e Pacific Ocean. At that  
aru" was located at about  
s nearly 100 tons and 23

in fishing tuna they saw  
on several minutes later.  
which continued for ap-  
returned to their mother

have been injured by the  
ed to the Tokyo Univer-  
First National Hospital  
n both hospitals in May  
mber 1954. After being  
r as possible on annual  
of their occupations and  
ible to get them all to-

ed several times (MIYOSHI  
t al., 1956; MIYOSHI and  
HI and KUMATORI, 1964;

## ERIALS

ts intensity was greatest,  
eyes open. Their foot-  
allout.

each patient's behavior on the boat. The dose to each person differed depending on his behavior and the position of his cabin.

The integrated dose to the thyroid from  $^{131}\text{I}$  was inferred as about 20 to 120 rads from external countings of radioactivity in the thyroid region during the 4 to 7 weeks after the initial exposure.

Significant amounts of radioactivity were found in the urine samples collected at about 4 weeks after the explosion. The radioactivity decreased rapidly, namely at about 6 months post explosion the activity in the urine was barely detectable.

Radiochemical analysis of several organs from the fatal case showed that the radioactivity of these organs was clearly higher, when compared with the controls, although it was not extremely high (KIMURA et al., 1956).

After 8½ years and 10 years,  $^{137}\text{Cs}$ - and  $^{90}\text{Sr}$  content in the urine of the fishermen revealed no significant increase. At the same time the results of whole body counting showed no significant difference between fishermen and controls.

Thus, the external irradiation seemed to play an important role in early effects.

#### SYMPTOMS AND SIGNS IN THE EARLY STAGE

Figure 1 summarizes the symptoms and signs in the early stage.




Fig. 1 Symptoms and signs in the early stage.

#### HEMATOLOGY

##### 1. Leukocytes

The total number of leukocytes at about the 4th to 6th week after the explosion was a count of less than 2,000 in one case, the leukocyte level was between these minimum counts. Similar relationship was observed between the dose as shown in Fig. 2.

Fig. 2 Correlation between the estimated doses to the whole body and the minimal number of neutrophils.

In most cases a shift in the leukocyte count was observed in early stages, especially with the bone marrow damage, a considerable number of cases. Lymphopenia and eosinophilia was seen in some cases. Eosinophilia continued in some cases and one case still revealed eosinophilia.

##### 2. Erythrocytes

At the time of the explosion, anemia was observed in the early stages. The Price-Jones curve was normal at first, and then it corresponded to changes in the leukocyte count.

##### 3. Platelets, etc.

Platelet counts were normal in most of the fishermen. Most of the fishermen had normal platelet counts in general and received no special treatment. When these blood counts were compared with the normal range, they were within the normal range.

## HEMATOLOGICAL OBSERVATIONS

## 1. Leukocytes

The total number of leukocytes decreased gradually, showing minimum counts at about the 4th to 8th week after the exposure. Five cases revealed a count of less than 2,000/mm<sup>3</sup>, 13 less than 3,000, and 5 less than 4,000. In one case, the leukocyte level was depressed to 800. A correlation was found between these minimum counts and the doses of individual external irradiation. Similar relationship was found between minimum counts of neutrophils and the dose as shown in Figure 2.




Fig. 2 Correlation between the estimated doses to the whole body and the minimal number of neutrophils.

In most cases a shift to the left of the neutrophils was observed. Staff cells increased in early stage and then gradually decreased. However, in some cases, metamyelocytes and myelocytes appeared in peripheral blood especially with the begin of recovery. In a fatal case who died of liver damage, a considerable number of juvenile neutrophils were observed at the end stage. Lymphopenia was noted between the 2nd and 8th week in all cases. Leukocyte count began to increase after the 8th week. In many cases eosinophilia was remarkable, especially with the indication of recovery. Eosinophilia continued in some cases for several years without proof of parasites, and one case still revealed slight eosinophilia in the 1968 survey.

## 2. Erythrocytes

At the time of hospitalization a few patients were anemic. No reticulocyte was observed in them at the critical stage. Color index was higher than 1.0. The Price-Jones curves of erythrocyte diameter were displaced to the right of normal at first, and returned to almost normal after one year. These changes corresponded to changes of erythroblast diameters in bone marrow.

## 3. Platelets, etc.

Platelet counts decreased, reaching a minimum in the 4th to 7th week. Most of the fishermen revealed the values below 100,000/mm<sup>3</sup>. The results of other tests related to hemorrhage showed depression at the critical stage in general and recovered by the 10th week.

## 4. Comparison with Other Irradiated Cases

When these blood changes are compared with the changes of other irradiated

use to each person differed de- his cabin.

<sup>131</sup>I was inferred as about 20 to 100 in the thyroid region during

found in the urine samples. The radioactivity decreased from the activity in the urine was

from the fatal case showed that other, when compared with the (KIMURA et al., 1956).

Sr content in the urine of the at the same time the results of difference between fishermen and

lay an impotant role in early

## EARLY STAGE

signs in the early stage.

Fig. 1 Symptoms and signs in the early stage.

subjects, *i. e.* Hiroshima cases, Marshallese, and cases exposed by reactor accidents (HEMPELMAN et al., 1952; AMANO, 1953; CRONKITE et al., 1956; HASTERLIK and MARINELLI, 1956; GUSKOVA and BAISSOLOV, 1956; ANDREWS et al., 1961; JAMMET, 1961), it is noted that all these cases similarly revealed the minimal value at about one month after exposure.

#### 5. Bone Marrow

The bone marrow was aplastic at the critical stage in severe cases, which showed remaining and proliferation of plasma cells and reticulum cells. Figure 3 shows the bone marrow of one of the severe cases. Comparing with normal




Fig. 3 Bone marrow of one of severe cases at 5th week.

bone marrow a lack of bone marrow cells was observed. With the beginning of recovery, aplastic marrow changed to hypoplastic and then turned into a type of maturation arrest. In some cases, recovery was not complete even after one year. In less severe cases, the bone marrow was not aplastic but hypoplastic or in a type of maturation arrest even at the critical stage.

#### 6. Morphological and Functional Changes

Several morphological abnormalities were observed: nuclear debris of lymphocytes, binucleated lymphocytes, vacuoles of neutrophils, toxic granulations of neutrophils, degenerative changes of blood cells, mitotic abnormalities of erythroblasts, etc.

Motility and phagocytosis of neutrophils decreased remarkably. After one year, in some cases these functions were still lower than in normal persons.

### FOLLOW-UP STUDIES

As above mentioned 16 to 18 fishermen have been examined on an annual basis.

Figure 4 shows the changes of the cumulative distribution curves of leukocytes. The curve of critical stage appeared displaced far to the left of the normal curve, and gradually approached normal values.

However, the curve of the 6th year was still displaced slightly to the left

Fig. 4 Leukocytes of Biki (J) (cumulative).

Fig. 5 Changes in leukocytes, neutrophils and

of normal Japan the curve approached the cumulative

However, the changes of are representative


A small increase of erythroblasts 10 years.

Cytogenetic results are summarized

1. The

and cases exposed by reactor ac-  
 3; CRONKITE et al., 1956; HASTERLIK  
 LOV, 1956; ANDREWS et al., 1961;  
 es similarly revealed the minimal

critical stage in severe cases, which  
 cells and reticulum cells. Figure  
 cases. Comparing with normal



s at 5th week.

observed. With the begin-  
 aplastic and then turned  
 recovery was not complete  
 marrow was not aplastic  
 n at the critical stage.

ed: nuclear debris of  
 phils, toxic granulations  
 mitotic abnormalities of

emarkably. After one  
 n in normal persons.

examined on an an-

ion curves of leuko-  
 to the left of the

slightly to the left

Fig. 4 Leukocytes of Bikini patients (J) (cumulative).





Fig. 5 Changes in the number of leukocytes, neutrophils and lymphocytes.

of normal Japanese. In the case of erythrocytes, 2 years after the exposure, the curve approximately came back to normal. A similar tendency is seen in the cumulative distribution curves of platelets.

However, in a few cases slight neutropenia is still observed. Figure 5 shows the changes of total leukocytes, neutrophils, and lymphocytes of 2 cases which are representative for severe injury.

A small increase of "mitotically connected abnormalities", i. e. karyomeres of erythroblasts, were observed in bone marrow smears of a few cases after 10 years.

Cytogenetical studies have been performed since several years. The results are summarized as follows (ISHIHARA and KUMATORI 1965; 1967; 1969).

1. The frequency of aneuploid cells was 2~3 per cent, which was not

**Table 2** Chromosome abnormalities in peripheral lymphocytes of the fishermen, data from 1965 to 1967.

72-h culture:					
Cases	No. of cases examined	No. of cells analyzed	Aneuploid cells %	Stable cells % ( / )*	unstable cells % ( / )*
Control	10	693	2.88	0.14 (1/10)	0.14 (1/10)
Fishermen 1965	13	881	2.05	2.28 (9/13)	0.46 (3/13)
Fishermen 1966	15	1102	2.90	3.45 (14/15)	0.45 (5/15)
Fishermen 1967	16	1248	2.48	2.88 (15/16)	0.56 (6/16)
48-h culture:					
Fishermen 1966	10	750	2.27	1.20 (6/10)	0.67 (4/10)
Fishermen 1967	12	950	2.00	2.11 (9/12)	0.74 (6/12)

\* Number of cases showing the abnormal cells per total number of cases examined.

so high (Table 2).

2. Unstable abnormalities such as dicentrics, etc. were observed (Table 2).
3. The frequency of stable abnormalities was remarkably higher than in controls (Table 2).
4. The detailed analysis of stable cells revealed that stable abnormalities were divided following 3 types:
  - a. stable cells with balanced chromosome constitution... 67 per cent
  - b. stable cells with deletion of a chromosome... 29 per cent
  - c. stable cells with excessive chromosome material... 4 per cent
5. In 3 cases small chromosomes which are similar to the Ph<sup>1</sup> chromosome were found.
6. The examined fishermen were classified in 3 groups according to the degree of the injuries indicated by the minimum values of neutrophils shown shortly after the exposure. It was found that the mean values of stable and of unstable cells were higher in proportion to the extent of the initial damage. Minimum values of neutrophils roughly corresponded to the estimated external radiation dose.
7. Chromosome abnormalities in bone marrow were observed in 5 cases. All of these abnormalities were limited to stable abnormalities. While stable cells in the peripheral lymphocytes showed various karyotypes of their own, these in the bone marrow were limited to only a few karyotypes.

### SPERMATOGENESIS

As a generative tissue, spermatogenesis has many similarities with hematopoiesis. Therefore, the changes of number of spermatozoa is described for reference. The number of spermatozoa decreased, and minimum numbers were obtained in 16 cases about 8 months after the exposure. Lowering of mobility

and morphological abnormalities of recovery were observed. First signs of regeneration were observed, with following:

The hematological findings of 1954 were reported with a dose soon after the accident. Generally speaking, decisions should be decided on the basis of the findings. From this point of view, it is necessary to decide the therapy necessary to detect the abnormalities observed in hematological

- AMANO, S. (1953): *Gakujutsu Shi*
- ANDREWS, G. A.: *S*  
Diagnosis and  
Organization.
- CRONKITE, E. P.:  
Ionizing Rad  
Washington.
- CUSKOVA, A. K. a  
Conference  
New York.
- HASTERLIK, R. J.:  
Conference  
New York.
- HEMPELMAN, L.:  
Medicine, :
- ISHIHARA, T. a  
ISHIHARA, T. a  
North Ho
- ISHIHARA, T. a  
242.
- JAMMET, H. P.:  
World H.
- KIKUCHI, T.:  
stitute f
- KIMURA, K.:  
Influenc  
the Pro



ocytes of the fishermen,

Stable cells % ( / )*	unstable cells % ( / )*
0.14 (1/10)	0.14 (1/10)
2.28 (9/13)	0.46 (3/13)
3.45 (14/15)	0.45 (5/15)
2.98 (15/16)	0.56 (6/16)
<hr/>	
1.20 (6/10)	0.67 (4/10)
2.11 (9/12)	0.74 (6/12)

Number of cases examined.

were observed (Table 2).  
Remarkably higher than

that stable abnormalities

... 67 per cent

29 per cent

1... 4 per cent

to the Ph<sup>1</sup> chromo-

groups according to the

values of neutrophils

mean values of stable

extent of the initial

bonded to the estimat-

observed in 5 cases.

abilities. While stable

otypes of their own,

aryotypes.

ilarities with hema-

zoa is described for

imum numbers were

owering of mobility

and morphological abnormalities of spermatozoa were also observed. Indications of recovery were observed in some cases after about one year, but mostly first signs of regeneration did not show earlier than about 2 years after exposure, with following production of children.

#### SUMMARY

The hematological changes of Japanese fishermen exposed to fallout in 1954 were reported with other findings.

Generally speaking, it is difficult to estimate correctly the exposed radiation dose soon after the accident. Therefore, treatment of heavily irradiated persons should be decided by the daily appraisal of clinical and laboratory findings. From this point, hematological changes are an important information to decide the therapy which should be given to heavily irradiated subjects.

As to the Japanese fishermen, it seems to be very important as well as necessary to detect the significance of subtle changes such as they are still observed in hematological and cytogenetical examinations.

#### REFERENCES

- AMANO, S. (1953): Genshi Bakudan Saigai Chosa Hokokushu. II, 914. Nihon Gakujutsu Shinko Kai, Tokyo.
- ANDREWS, G. A., SITTERSON, B. W., KRETCHMAR, A. L. and BRUCER, M. (1961): Diagnosis and Treatment of Acute Radiation Injury, 27, World Health Organization, Geneva.
- CRONKITE, E. P., BOND, V. P. and DUNHAM, C. H. (ed.) (1956): Some Effects of Ionizing Radiation on Human Beings. U.S. Government Printing Office, Washington, D. C.
- GUSKOVA, A. K. and BAISSOLOV, G. D. (1956): Proceedings of the International Conference on Peaceful Uses of Atomic Energy, 11, 35, United Nations, New York.
- HASTERLIK, R. J. and MARINELLI, L. D. (1956): Proceedings of the International Conference on Peaceful Uses of Atomic Energy, 11, 25, United Nations, New York.
- HEMPELMAN, L. H., LISCO, H. and HOFFMAN, J. G. (1952): Annals of Internal Medicine, 36 279.
- ISHIHARA, T. and KUMATORI, T. (1965): Acta Haematologica Japonica, 28, 291.
- ISHIHARA, T. and KUMATORI, T. (1967): Human Radiation Cytogenetics, 144, North Holland Publishing Co., Amsterdam.
- ISHIHARA, T. and KUMATORI, T. (1969): Japan Journal of Genetics, 44, Suppl. 1, 242.
- JAMMET, H. P. (1961): Diagnosis and Treatment of Acute Radiation Injury, 83, World Health Organization, Geneva.
- KIKUCHI, T., WAKISAKA, G., AKAGI, H. and GOTO, H. (1954): Bulletin of the Institute for Chemical Research, 4, Kyoto University, Kyoto.
- KIMURA, K., MINAMI, E., HONDA, M. et al. (1956): Research in the Effects and Influences of the Nuclear Bomb Test Explosions, I, 497, Japan Society for the Promotion of Science, Tokyo.

- KIMURA, K., IKEDA, N., KIMURA, K., KAWANISHI, H. and KIMURA, M. (1956): *ibid.* I, 521.
- KOYAMA, Y., KUMATORI, T. et al. (1955): *Iryo*, 9, 1.
- KUMATORI, T. and MIYOSHI, K. (1963): *Diagnosis and Treatment of Radioactive Poisoning*, 253, International Atomic Energy Agency, Vienna.
- KUMATORI, T., ISHIHARA, T., UEDA, T. and MIYOSHI, K. (1965): *Medical Survey of Japanese Exposed to Fallout Radiation in 1954—A Report after 10 Years*, National Institute of Radiological Sciences, Chiba.
- MIKAMO, Y., MIYOSHI, K., SHIMIZU, K., ISHIKAWA, K., KURIYAMA, S., KOYAMA, Y. and KUMATORI, T. (1956): *Research in the Effects and Influences of the Nuclear Bomb Test Explosions*, II, 1313, Japan Society for the Promotion of Science, Tokyo.
- MIYOSHI, K. and KUMATORI, T. (1955): *Acta Haematologica Japonica*, 18, 379.
- MIYOSHI, K. and KUMATORI, T. (1962): *Proceedings of 8th International Congress of Hematology*, 1, 29, Pan-Pacific Press, Tokyo.
- MIYOSHI, K. and KUMATORI, T. (1964): *Nihon Ketsuekigaku Zensho*, 3, 660, Maruzen Co., Tokyo.

## DISCUSSION

Dr. CLIFTON (U.S.A.): Concerning the one fatality, would you care to comment on the nature of the liver damage, and its relationship to external or internal radiation exposure?

Dr. KUMATORI (Japan): Since this fatal case was anemic and revealed bone marrow aplasia, he received blood transfusions. Therefore, serum hepatitis can't be ignored. However, at the same time the existence of radiation-induced liver damage should be considered. At present it is difficult to decide the cause of his liver damage.

We should consider the probability of the radiation-induced liver damage, because in some publications we can see a similar histological picture of the liver of a patient who was irradiated therapeutically.

Dr. SANG (Korea): I thank you Dr. KUMATORI, I would like to ask you something about hemorrhagic tendencies on these cases, Would you tell me how long did hemorrhagic tendencies continue after the irradiation?

Dr. KUMATORI: In my cases the hemorrhagic tendencies were not so severe. The prolongation of the bleeding time was observed in a few cases. You can see normal bleeding time in

most cases. In our cases, we used DUKE's method. The bleeding time is normally 3 minutes, and in 1 or 2 severe cases the bleeding time at the critical stage was as long as about 10 minutes. I think this came back to normal at about 10 weeks or so after the exposures, though I have no correct data here.

Dr. SANG: And how about the fibrinogen concentration?

Dr. KUMATORI: Fibrinogen concentration was almost normal from the beginning of the examination.

Dr. TUBIANA (France): I would like to ask Prof. KUMATORI if he has an idea of how uniform was the dose delivered to your fishermen. I ask you this question because we have a rather large experience of total-body irradiation for mostly kidney transplants in human patients, and there are 2 main differences between your results and the one we have also. The first one is that aplasia of the blood occurs much earlier in our patients, and the rigidity also occurs much earlier. The 2nd one is that the minimum number of leukocytes is much smaller in our cases, being of the order 1 to 200 leukocytes per cubic millimeter, after a dose of the

order of 350 R to 40 and LALANNE, C. M., *radiologie*, 6, 561, (1963) and *Ionizing Radiation on Hemopoietic tissue*, *Int. At. Energy Agency* (1967)).

We have compared the curves of Yugo cases, and here again same difference, and attention of these data and the difference between cases and the patients was that patients the dose with a maximum value minus 4 per cent, and of accidents the un-

Dr. KUMATORI: In my opinion. In my c

order of 350 R to 400 R (TUBIANA, M. and LALANNE, C. M., *Annales de Radiologie*, 6, 561, (1963) and in *Effects of Ionizing Radiation on the haematopoietic tissue*, Int. At. En. Ag., Vienna (1967)).

We have compared our curves with the curves of Yugoslavian accident cases, and here again we found the same difference, and our interpretation of these data was that probably the difference between the irradiated cases and the purposely irradiated patients the dose was very uniform, with a maximum variation of plus or minus 4 per cent, whereas in the case of accidents the uniformity was much less, and of the order of 30 per cent.

Dr. KUMATORI: I agree with your opinion. In my cases, I think, the

main irradiation came from externally deposited sources--gamma irradiation. This is a type of uniform irradiation. And this irradiation played an important role in the hematological changes. The fallout material began to precipitate at 7:00 a.m. and continued to fall for about 5 hours. Therefore, the radiation dose to fishermen was at first small and then rapidly increased. As the fishermen began to clean up at about 12:00, the exposure rate decreased thereafter. They were irradiated continuously for 2 weeks with decreasing dose rates. Aside from this external irradiation, internal exposure may have had some effects. But the internal irradiation seemed to have only a slight effect on the acute hematological changes.

A. M. (1956): *ibid.*

ent of Radioactive  
enna.

Medical Survey of  
ort after 10 Years,

S., KOYAMA, Y. and  
es of the Nuclear  
otion of Science,

onica, 18, 379.

ational Congress

Zensho, 3, 660,

cases, we used  
bleeding time is  
and in 1 or 2  
ling time at the  
ong as about 10  
s came back to  
eeks or so after  
gh I have no

about the fib-

nogen concent-  
rmal from the  
ination.

I would like  
if he has an  
was the dose  
ermen. I ask  
use we have a  
of total-body  
kidney trans-  
nts, and there  
between your  
we have also.  
plasia of the  
earlier in our  
ty also occurs  
one is that the  
leukocytes is  
ases, being of  
ukocytes per  
a dose of the