



BROOKHAVEN NATIONAL LABORATORY  
ASSOCIATED UNIVERSITIES, INC., UPTON, L.I., N.Y. 11973

410320

HEALTH PHYSICS & SAFETY DIVISION

TELEPHONE: (516) 345-4250

December 18, 1975

Mr. Tommy F. McCraw  
Division of Operational Safety  
U.S. Energy Research and Development  
Administration  
Washington, D. C. 20545

Dear Tommy:

Thanks a million for "bailing me out" on the paper. We got it off yesterday (I hope within the grace period for inclusion in the meeting proceedings). In any event, I'm enclosing some copies for your use or disposal.

Best wishes for the holiday season.

Sincerely yours,

A handwritten signature in cursive script, appearing to read 'Nat'.

N. A. Greenhouse

1k  
Encs.

*McCraw,*

DRAFT

December 5, 1975

MARSHALL ISLANDS RADIOLOGICAL FOLLOWUP

by

N. A. Greenhouse  
Health Physics and Safety Division  
Brookhaven National Laboratory

and

T. F. McCraw  
Division of Operational Safety  
U.S. Energy Research and Development Administration

ABSTRACT

By acceptance of this article, the publisher and/or recipient acknowledges the U.S. Government's right to retain a nonexclusive, royalty-free license in and to any copyright covering this paper.

Research carried out at Brookhaven National Laboratory under the auspices of the U.S. Energy Research and Development Administration.

## INTRODUCTION

The U. S. nuclear testing programs of the 1940s and 1950s had significant local environmental impacts on the coral atolls of Bikini and Enewetak in the Marshall Islands. The high level close in fallout made these atolls uninhabitable for many years. Fallout from the BRAVO event, which took place at Bikini in 1954, was inadvertently deposited on the nearby Atolls of Rongelap, Rongerik and Utirik. In all, some thirteen atolls in the northern Marshalls were probably affected to a greater or lesser extent by fallout from these nuclear tests. Of these, however, the most significant long-term radiological impact was for the test Atolls, Bikini and Enewetak, and for Rongelap Atoll.

In 1957, Rongelap was reoccupied by its original inhabitants who had been evacuated within about two days after BRAVO. During the past several years, definitive plans have been made to repatriate the original inhabitants of Bikini and Enewetak Atolls, and their families. It is hoped that their return can take place soon.

In order to identify radiological problems from residual radioactivity in the environment, and to provide a data base for dose predictions applicable to the returning populace, ERDA (and its predecessor, the AEC), has sponsored many radiological surveys in the Marshall Islands. These surveys began during test operations and have been conducted periodically up to the present time. Results of the surveys have been published in numerous reports and scientific journals. References 1 through 12 are published reports of AEC/ERDA supported surveys of these atolls. References 13 through 29 are a portion of the published reports on work with collected environmental samples supported by AEC/ERDA.

Evaluation of survey results for Bikini Atoll, the consideration of predicted exposures compared with applicable radiation standards, and the acknowledgement of the many benefits to the people if they could return,

led to the decision to clean up and rehabilitate that Atoll. The Department of Defense, Department of the Interior (DOI), and AEC (now ERDA) participated in a joint effort of cleanup and rehabilitation of Bikini Atoll starting in February of 1969. Cleanup was completed in the fall of that year. Agricultural rehabilitation and housing construction is being conducted by DOI.

The decision to return the Enewetakese to their atoll led to a comprehensive survey conducted at Enewetak in 1972-1973. (See reference 10) A regional survey planned for 1976 will provide base-line radiological data for future dose assessments throughout nearly all of the northern Marshall Islands which may have been affected by the testing program. Environmental evaluations at Rongelap and Utirik Atolls have been undertaken periodically in association with ERDA's medical evaluations program there over the past 20 years. (Nat, suggest we cite Bob C. references here.)

From all of these earlier surveys, it became apparent that periodic environmental monitoring and dose assessments must be made for Bikini, Enewetak, Rongelap and perhaps other atolls in the northern Marshall's to maintain a current radiological data base and to provide current information on individual and population doses. This followup monitoring is being performed by Brookhaven National Laboratory at the request of the Division of Operational Safety, ERDA.

#### RADIOLOGICAL CONCERNS

The primary radiological problems are the result of residual fission and activation products in the terrestrial environment. They have been identified by previous environmental surveys as follows:

1. External radiation levels significantly higher on some islands in an atoll compared to levels on lightly contaminated islands.
2. Fission and activation product radioactivity in certain terrestrial

food items now growing on islands of these atolls and the possibility that unacceptable levels of these radionuclides may appear in foods, plants and animals newly introduced into these atolls.

3. Radioactivity in the ground water, a possible source of drinking water and water for irrigation.
4. Plutonium and americium isotopes in the surface soil.

These factors are illustrated by data in the accompanying tables taken from previous radiological survey reports. (See Tables 1, 2, 3, and 4.)

In some cases, the predicted doses and dose commitments derived from survey information for Bikini and Enewetak Atolls approach or even exceed national and international radiation protection standards for certain living and dietary patterns. Corrective actions or restrictions must be placed on use of these Atolls and their resources to assure that the applicable radiation standards are not exceeded. Herein lies the primary justification for the continuing environmental followup surveys sponsored by ERDA.

#### ENVIRONMENTAL MONITORING

The most important sources of exposure to people living on Rongelap and to future residents of Bikini and Enewetak Atolls are from internal deposition of radioisotopes from certain elements in the human diet, and from the long-term occupancy of islands having external radiation dose rates higher than natural background. Aside from periodic re-evaluations to establish trends in external dose rate reduction, external radiation monitoring will assume less significance, compared to monitoring of the food chain, as time passes. At present, annual visits are being made to identify and collect representative samples of local diets for laboratory analysis and dose commitment updates. New locally grown food items are becoming available in small quantities on Bikini Island as a result of the

experimental agricultural practices of a small group of caretaker families living there. Neither Bikini Atoll, where radiological clean-up has been completed, nor Enewetak Atoll where cleanup has not yet begun, have a subsistence agriculture resource in being which is sufficient to support the anticipated populations which will one day live there ( though such crops are currently being developed or planned). As a result, some of the dietary items likely to have the higher radionuclides content, e. g. , pandanus and breadfruit, are not actual problems to date. They may or may not be of concern in the future as the plantings mature and the fruit becomes available in quantity. Thus, the diets of people living in these two atolls are expected to change over the coming years reflecting the relative influences of imported and locally grown food items. Allowance has been made for this in development of radiation dose estimates. Experimental studies at Enewetak may yield techniques to interrupt or break the recycling of radionuclides through the vegetation, soil, and ground water systems, and thereby reduce the radioactivity content of some important dietary items. All of the aforementioned factors will necessitate continuing monitoring of the diet for many years. Periodic sampling and analysis of soil and ground water will be necessary in order to establish trends in the changes of radioactivity content of these media.

In the Northern Marshalls, drinking water is obtained primarily from rainwater catchments. While the radionuclide content of collected rainwater will not be zero, this source is not expected to contribute significantly to the radiation exposure picture for future Bikini, Enewetak, and Rongelap Atoll residents. However, rainwater which drains from the windward side of building rooftops may provide useful data on re-suspension of radioactivity in the soil. The collection of rainwater by

future Bikini and Enewetak residents is being facilitated by including gutters and water storage tanks in plans for houses and community structures. Some of the larger islands have fresh ground water located only a few feet below the surface. Analysis of this water for its radionuclide content has been limited to date and the capacity of this resource to serve the needs of island residents is not well defined. More study of this water is being supported by ERDA.

#### PERSONNEL MONITORING

Dose predictions for Bikini and Enewetak Atoll residents derived from environmental data have been deliberately conservative, and establish probable upper limits on doses to be expected for individuals. Reliable assessments of actual doses must be determined through personnel monitoring. External radiation dosimeters do not appear to be a practical means of personnel monitoring for individual external dose measurements, although certain individuals within given populations may be relied upon to wear them. A "lifestyle model" which includes estimates of occupancy factors for various locations in a given atoll has been coupled with environmental monitoring data to estimate average external radiation doses to individuals. This model will be revised as needed so that it closely approximates the actual lifestyle of the people.

The more important internal pathway can be monitored directly by conventional techniques of bioassay and whole-body counting of individuals. A portable shadow shield whole-body counter has been constructed and mounted in a shipboard trailer for use in the Marshall Islands. It is capable of quantitative detection of very small quantities of certain radionuclides in the body such as  $^{137}\text{Cs}$  and  $^{60}\text{Co}$ , the primary environmental gamma emitters at Bikini, Enewetak, and Rongelap Atolls. The system

clearly identifies individuals in the Rongelap population who are not following the recommended dietary restrictions on eating coconut crabs from certain locations. Body burdens of  $^{90}\text{Sr}$ ,  $^{90}\text{Y}$ ,  $^{239, 240}\text{Pu}$  and  $^{241}\text{Am}$  are estimated by the radiochemical analysis of urine samples. Urine sample collections and whole-body counting will be performed every 1 to 2 years at Bikini and Enewetak Atolls when the people return, and every 2 to 3 years at Rongelap Atoll until the results warrant less frequent measurement intervals.

### SUMMARY

Marshall Islands Radiological Followup has consisted of intensive environmental studies at Bikini, Enewetak, and Rongelap Atolls to gather radiological data on the external radiation environment and on radioactivity in food chains. Radiation and radioactivity levels in these atolls are being reduced with time. These changes are monitored in annual or bi-annual environmental surveys. Updated information is used to make conservative estimates of population doses and dose commitments. When people have returned, actual internal doses to individuals are determined for whole-body counting and bioassay data. These results are combined with environmental data on the external radiation environment to complete the total dose assessment picture.



## REFERENCES

- Dunning, G. M. - Radioactive Contamination of Certain Areas in the Pacific Ocean from Nuclear Tests, A Summary of the Data from the Radiological Surveys and Medical Examinations, USAEC Report, August 1957.
- Held, E. E. - Gamma Dose Rates at Rongelap Atoll, 1954-1963, USAEC Report, UWFL-91, May 1965.
- Donaldson, L. R., et al, Bikini-Eniwetok Studies, 1964, Part I and II, University of Washington, Laboratory of Radiation Biology, UWFL-93, September 15, 1966.
- Beck, H. L., Bennett, B. G. and McCraw, T. F. - External Radiation Levels on Bikini Atoll, May 1967, USAEC Report HASL-190, December 1967.
- Bennett, B. G. and Beck, H. L., -External Radiation on Bikini Atoll, Nature, Vol. 223, No. 5209, August 30, 1969.
- Held, E. E. - Radiological Resurvey of Animals, Soils and Ground Water at Bikini Atoll, 1969-1970, University of Washington, College of Fisheries, Laboratory of Radiation Ecology, NVO-269-8 (Revision 1), February 1971.
- Smith, A. E. and Moore, W. E. - Report of Radiological Cleanup of Bikini Atoll, Office of Dose Assessment and Systems Analysis, Western Environmental Research Laboratory, Environmental Protection Agency, SWRHL-111r, January 1972.
- McCraw, T. F. and Lynch, O. D. T. Jr. - Exposure Rate Reduction on Bikini Island Due to Concrete Dwellings, USAEC Report, WASH-1273, June 1973.
- McCraw, T. F. - Levels of Environmental Radioactivity in Bikini Atoll, USAEC Report, WASH-1289.
- Enewetak Radiological Survey, Vol. I, II, and III, USAEC Nevada Operations Office Report NVO-140, October 1973.
- Lynch, O. D. T. Jr., McCraw, T. F., Nelson, V. A., and Moore, W. E. - Radiological Resurvey of Food, Soil, Air and Ground Water at Bikini Atoll, 1972, USERDA Report ERDA-34, February 1975.
- Gudixsen, P. H., and Robison, W. L. - Preliminary External-Dose Estimates for Future Bikini Atoll Inhabitants, Lawrence Livermore Laboratory Report, Preliminary UCRL-51879, August 6, 1975.

- Palumbo, R. F. and Lowman, F. G. - The Occurrence of Antimony-125, Europium-155, Iron-55, and Other Radionuclides in Rongelap Atoll Soil, USAEC Report UWFL-56, April 7, 1958.
- Lowman, F. G. - Marinebiological Investigations at the Eniwetok Test Site. Reprinted from Disposal of Radioactive Wastes, International Atomic Energy Agency Report, 1960.
- Walker, R. B., Held, E. E. and Gessel, S. P. - Radiocesium in Plants Grown on Rongelap Atoll Soils, Recent Advances in Botany, University of Toronto Press - 1961.
- Cole, D. W., Gessel, S. P. and Held, E. E. - Tension Lysimeter Studies of Ion and Moisture Movement in Glacial Till and Coral Atoll Soils, Soil Science Society of America Proceedings, Vol. 25, No. 4, July-August 1961.
- Lowman, F. G. and Palumbo, R. F. - Occurrence of Bismuth-207 at Eniwetok Atoll, Nature, Vol. 193, No. 4817, February 24, 1962.
- Chakravarti, D. and Held, E. E. - Chemical and Radiochemical Composition of the Rongelapese Diet, Journal of Food Science, Vol. 28, No. 2, 1963.
- Kenady, R. M. - The Soils of Rongelap Atoll, Marshall Islands, University of Washington - 1962.
- Held, E. E. - Qualitative Distribution of Radionuclides at Rongelap Atoll, Radioecology - 1963.
- Held, E. E., Gessel, S. P., Matson, L. J. and Billings, R. F. - Autoradiography of Sectioned Soil Cores, Radioisotope Sample Measurement Techniques in Medicine and Biology, International Atomic Energy Agency, Vienna, 1965.
- Held, E. E., Gessel, S. P., and Walker, R. B. - Atoll Soil Types in Relation to the Distribution of Fallout Radionuclides, USAEC Report, UWFL-92, August 1965.
- Welander, S. D. - Distribution of Radionuclides in the Environment of Eniwetok and Bikini Atolls, August 1964, Symposium on Radioecology, Proceedings of Second National Symposium, Ann Arbor, Michigan, May 15-17, 1967, CONF-670503.

Beasley, T. M. and Held, E. E. - Nickel-63 in Marine and Terrestrial Biota, Soil, and Sediment, Science, Vol. 164, June 6, 1969.

Beasley, T. M. and Held, E. E. - Silver-108m in Biota and Sediments at Bikini and Eniwetok Atolls, Nature, Vol. 230, No. 5294, April 16, 1971.

Schell, W. R. and Yang, A. I. C. - Long-lived Radionuclides Produced at Bikini and Eniwetok Atolls, Laboratory of Radiation Ecology, College of Fisheries, University of Washington, RLO-2225-T18-3, April 30, 1973.

Schell, W. R. - Studies of Concentrations of Unreported Long-lived Radionuclides in Biota and Ocean Sediments at Bikini and Eniwetok Atolls, Laboratory of Radiation Ecology, College of Fisheries, University of Washington, RLO-2225-T18-4, May 1, 1973.

Noskin, V. E., Wong, K. M., Eagle, R. J. and Gatrousis, C. - Lawrence Livermore Laboratory Report, UCRL-51612, June 26, 1974.

Nevissi, A., Schell, W. R. and Nelson, V. A. - Plutonium and Americium in Soils of Bikini Atoll, IAEA/ERDA Symposium on Transuranium Nuclides in the Environment, San Francisco, Calif., November 17-21, 1975, RLO-2225-T18-17, IAEA-SM-199/62.

Table 1

Gamma Radiation Rates in Bikini Atoll

| Island                  | Exposure Rate Range<br>mR/hr | Major Contributors   |
|-------------------------|------------------------------|--|
| Bikini                  | .010-.120                    | <sup>137</sup> Cs  |
| Weathered Areas         | .010-.030                    |  |
| Close-to-Shore          | .020-.040                    |  |
| Island Center           | .050-.080                    |  |
| Hot Spots               | .080-.120+                   |  |
| Eneu                    | .002-.010                    | <sup>137</sup> Cs  |
| Nam                     | .010-.330                    | <sup>60</sup> Co, <sup>137</sup> Cs                        |
| Outer Edge              | .010-.030                    |  |
| Island Center           | .015-.150                    |  |
| N.E. Corner             | .110-.330                    |  |
| Bokantuak, Iomelan,     | .003-.010                    | * ?  |
| Rojkere, Eonjebi        |                              |  |
| Aerokoj-Eneman Complex: |                              |  |
| Aerokoj, Aerokojlol     | .001-.010                    | * ?  |
| Bikdrin, Lele           | .006-.010                    | * ?  |
| Eneman                  | .001-.570                    | <sup>60</sup> Co, <sup>125</sup> Sb,<br><sup>102m</sup> Rh |
| East Eneman             | .001-.010                    |  |
| West Eneman             | .020-.570                    |  |
| Enidrik                 | .003-.235                    | <sup>60</sup> Co, <sup>125</sup> Sb,<br><sup>102m</sup> Rh |
| East Enidrik            | .003-.030                    |  |
| West Enidrik            | .010-.235                    |  |
| Lukoj                   | .060-.200                    | <sup>60</sup> Co, <sup>125</sup> Sb,<br><sup>102m</sup> Rh |
| Jelete                  | .060-.130                    | * ?  |
| Oroken                  | .015-.045                    | * ?  |
| Bokaetoktok             | .010-.035                    | * -  |
| Bokdrolul               | .020-.050                    |  |
| Bokbata                 | .010-.030                    | <sup>60</sup> Co, <sup>137</sup> Cs                        |
| Aomen-Iroi Complex:     |                              |  |
| Aomen                   | .005-.020                    | * -  |
| Lomilik                 | .020-.330                    | <sup>60</sup> Co, <sup>125</sup> Sb                        |
| Oodik, Iroi             | .010-.040                    | * ?  |

\*No soil sample or field spectra measurements.

*see reference 9.*

percent  
10 to  
sites de-  
signifi-  
Am, and

Figure 2,

ii, the  
the gamma  
measure-  
essential-  
Transect  
icantly

sa for  
on popula-  
to the ex-  
and conc-  
center of  
, was about  
spectro-  
made be-  
cular area  
ion levels

vegetation  
to type,  
ential  
is small

a has little  
ring of  
rub plants  
ed two  
be about

any buildings  
vegetation  
ad a road  
rements  
ore on the  
rements  
re placed

Table 2. Concentrations of  $^{40}\text{K}$ ,  $^{60}\text{Co}$ ,  $^{90}\text{Sr}$ , and  $^{137}\text{Cs}$  in Food Plants Collected on Bikini Island in November, 1971, and March, 1972.

| Plant Type | Tissue          | Collection Date | pCi/g, dry <sup>a</sup> |                  |                   |                  |
|------------|-----------------|-----------------|-------------------------|------------------|-------------------|------------------|
|            |                 |                 | $^{40}\text{K}$         | $^{60}\text{Co}$ | $^{137}\text{Cs}$ | $^{90}\text{Sr}$ |
| Lettuce    | Entire          | March, 1972     | 7.1+6.1                 | NS <sup>b</sup>  | 1860+10           | 320 +25          |
| Papaya #1  | Seeds           | March, 1972     | 17 +2                   | 0.13+0.12        | 140+ 1            | NA <sup>c</sup>  |
| "          | Fruit           | " "             | 31 +3                   | NS               | 160+ 1            | 72 + 3           |
| Papaya #2  | Seeds           | March, 1972     | 14 +4                   | NS               | 270+ 2            | NA               |
| "          | Fruit           | " "             | 23, +3                  | NS               | 290+ 1            | 69 + 4           |
| Pandanus   | Fruit (edible)  | Nov. 1971       | 7.5+0.6                 | 0.05+0.03        | 82+ 0.4           | 56 + 0.6         |
| "          | Fruit (fibrous) | " "             | 12 +2                   | NS               | 200+ 1            | 220 + 2.2        |
| "          | Leaves          | " "             | 3.4+1.5                 | NS               | 71+ 0.4           | 190 + 1.9        |
| Coconut #1 | Meat            | Nov. 1971       | 1.6+1.3                 | NS               | 93+ 0.5           | NA               |
| "          | Milk            | " "             | 3.9+1.9                 | NS               | 110+ 0.7          | NA               |
| Coconut #2 | Meat            | Nov. 1971       | 4.4+2.1                 | NS               | 110+ 1            | 0.88+ 0.04       |
| "          | Milk            | " "             | 4.3+2.5                 | NS               | 100+ 1            | <.22             |
| Coconut #3 | Meat            | Nov. 1971       | 11 +4                   | NS               | 147+ 1            | NA               |
| Coconut #4 | Meat            | Nov. 1971       | 2.5+2.0                 | NS               | 100+ 0.8          | NA               |
| "          | Milk            | " "             | 3.0+1.8                 | NS               | 77+ 0.6           | NA               |
| Coconut #5 | Meat            | Nov. 1971       | 15 +6                   | NS               | 270+ 2            | NA               |
| "          | Milk            | " "             | 2.1+1.3                 | NS               | 33+ 0.3           | NA               |
| Coconut    | Fronds (old)    | Nov. 1971       | 7.0+5.0                 | NS               | 310+ 3            | NA               |
| "          | Fronds (new)    | " "             | 14 +5                   | NS               | 220+ 2            | NA               |

a. The error terms for  $^{40}\text{K}$ ,  $^{60}\text{Co}$ , and  $^{137}\text{Cs}$  are two-sigma, propagated, counting errors. The errors for  $^{90}\text{Sr}$  are one-sigma, propagated, counting errors.

b. NS = Not significant. The net sample count is less than the two-sigma, propagated, counting error.

c. NA = Not analyzed.

\* See reference 11.

3  
 Table 14. Some Radionuclides in Water Samples Collected with a Large-Volume Filter-Sorption Bed from Bikini Atoll, May 1972

| Collection Location                            | Fraction                 | Liters Sampled | Radionuclide Concentration in pCi/m <sup>3</sup> <sup>a</sup> |                   |                   |                   |                   |
|--|--------------------------|----------------|---|-------------------|-------------------|-------------------|-------------------|
|  |                          |                | <sup>60</sup> Co  | <sup>137</sup> Cs | <sup>155</sup> Eu | <sup>207</sup> Bi | <sup>241</sup> Am |
| Bravo Crater (Bottom)                          | Particulate <sup>b</sup> | 3785           | 51 +3   | 12 +1             | 97 +4             | 27 + 2            | 70 +5             |
|  | Soluble <sup>c</sup>     | "              | 28 +8   | <14               | <20               | 160 +11           | <30               |
| Bravo Crater (Surface)                         | Particulate              | 3785           | 6.5+1.4   | NS <sup>d</sup>   | 9.6+1.3           | 0.5+ 0.3          | 8.2+1.4           |
|  | Soluble                  | "              | <10   | <6                | <16               | <12               | <22               |
| Bokdrolul Pass (Ebb Tide)                      | Particulate              | 4088           | 6.0+7.3   | NS                | 6.4+1.2           | NS                | 5.4+1.1           |
|  | Soluble                  | "              | 1.4+0.6   | <1.0              | <2.3              | 3.0+ 0.5          | <2.9              |
| Bokdrolul Pass (Flood Tide)                    | Particulate              | 4921           | 2.1+0.7   | NS                | 1.5+0.8           | NS                | 1.8+0.9           |
|  | Soluble                  | "              | NS  | NS                | NS                | 2.5+ 1.7          | NS                |
| Bikini Island (Seaward Reef)                   | Particulate              | 1620           | 1.5+1.5   | NS                | NS                | 2.0+ 0.8          | NS                |
|  | Soluble                  | "              | 6.2+5.4   | NS                | NS                | NS                | NS                |
| Bikini Island (Lagoon)                         | Particulate              | 2271           | 5.6+1.0   | NS                | NS                | 0.76+0.38         | 1.1+1.1           |
|  | Soluble                  | "              | 9.2+6.6   | NS                | NS                | NS                | NS                |
| Ocean between Bikini and Eniwetok <sup>e</sup> | Particulate              | 4898           | NS  | 7.7+1.0           | NS                | NS                | NS                |
|  | Soluble                  | "              | NS  | NS                | NS                | NS                | NS                |
| Bikini Island (Freshwater well)                | Particulate              | 1893           | 21 +3   | 21 +1             | 14 +3.7           | NS                | NS                |
|  | Soluble                  | "              | 54 +9   | 990 +60           | <2                | <7                | 34 +0             |

- a. Errors are two-sigma, propagated, counting errors.
- b. Particulate - that portion retained by the 0.3μ filter.
- c. Soluble - that portion which passes through the 0.3μ filter and is sorbed by the Al<sub>2</sub>O<sub>3</sub> beds.
- d. NS = not significant. The net sample count is less than the two-sigma, propagated counting error.
- e. This sample was collected over a six-hour period between the following positions: 11°29'5" by 164°58'0"E to 11°24'5"N by 164°18'0"E.

4. See reference 11.

Table 1.  $^{239,240}\text{Pu}$ ,  $^{238}\text{Pu}$  and  $^{241}\text{Am}$  in surface soil samples collected at Bikini Atoll in 1972, activities in  $\text{pCi/g} \pm 1\sigma$ .

| Location  | $^{239,240}\text{Pu}$ | $^{238}\text{Pu}$ | $^{241}\text{Am}$ | $^{239,240}\text{Pu}/^{238}\text{Pu}$ | $^{239,240}\text{Pu}/^{241}\text{Am}$ |
|---|-----------------------|-------------------|-------------------|---------------------------------------|---------------------------------------|
| Isl-29 Ourukaen                                     | 16.3 ± 0.5            | 5.7 ± 0.2         | 3.6 ± 0.3         | 2.86                                  | 4.53                                  |
| Isl-30 Bokoatokutoku                                | 15.1 ± 0.7            | 3.9 ± 0.3         | 4.3 ± 0.42        | 3.87                                  | 3.51                                  |
| Isl-30 Rokoatokutoku<br>Pisonia Grove               | 22.2 ± 0.5            | 6.7 ± 0.2         | 7.0 ± 0.30        | 3.31                                  | 3.17                                  |
| Roro Bokororyuru<br>Isl-31 down center of island    | 36.4 ± 2.8            | 7.2 ± 0.8         | 13.0 ± 1.1        | 5.05                                  | 2.75                                  |
| Namu West end + 150 yds                             | 24.0 ± 0.1            | 0.28 ± 0.02       | 14.0 ± 0.4        | 85.7                                  | 1.68                                  |
| Namu 200 yds SW of Bunker<br>300 yds E of West tip  | 20.1 ± 0.3            | 0.24 ± 0.02       | 11.0 ± 0.7        | 83.8                                  | 1.76                                  |
| Namu Top of bunker center of<br>island              | 22.9 ± 0.7.           | 0.31 ± 0.04       | 15.0 ± 0.5        | 73.9                                  | 1.57                                  |
| Namu 200 yds E of bunker<br>Center of island        | 17.4 ± 0.6            | 0.57 ± 0.11       | 10.0 ± 0.5        | 30.5                                  | 1.68                                  |
| Bikini Row 24 Center BL to<br>1st BLN               | 3.3 ± 0.1             | 0.45 ± 0.04       | 2.2 ± 0.3         | 7.33                                  | 1.50                                  |
| Bikini N. corner of Ctr. BL<br>and Lagoon Beach Rd. | 3.41 ± 0.36           | N.R.*             | .87 ± 0.12        | -                                     | 4.01                                  |
| Bikini Row 34 Center BL to<br>1st BL                | 3.0 ± 0.2             | 0.06 ± 0.04       | 2.1 ± 0.2         | 50.0                                  | 1.42                                  |
| Bikini Row 38 2nd BLN to<br>Lagoon Beach Rd.        | 2.5 ± 0.2             | 0.07 ± 0.02       | 1.2 ± 0.2         | 35.7                                  | 2.08                                  |

Table 1. (cont.)

| Location  | $^{239,240}\text{Pu}$ | $^{238}\text{Pu}$ | $^{241}\text{Am}$ | $^{239,240}\text{Pu}/^{238}\text{Pu}$ | $^{239,240}\text{Pu}/^{241}\text{Am}$ |
|---|-----------------------|-------------------|-------------------|---------------------------------------|---------------------------------------|
| Bikini Row 25 or 26 Sandpile Sample. 100 yds S of 2nd BLN | $0.50 \pm 0.05$       | N.S.**            | -                 | -                                     | -                                     |
| Bikini Row 34 Ctr. BL to 1st BLS                          | $10.8 \pm 0.4$        | N.R.              | $3.3 \pm 0.3$     | -                                     | 3.27                                  |
| Bikini Row 24 Ctr. BL to 1st BLS                          | $13.2 \pm 0.3$        | N.R.              | $8.4 \pm 0.55$    | -                                     | 1.58                                  |
| Bikini Row 38 2nd BLN to 1st BLN                          | $13.1 \pm 0.6$        | $0.74 \pm 0.10$   | $7.7 \pm 0.2$     | 17.7                                  | 1.70                                  |
| Bikini Row 24 1st BLN to Lagoon Beach Rd.                 | $9.3 \pm 0.4$         | $0.39 \pm 0.07$   | $4.1 \pm 0.2$     | 23.8                                  | 2.27                                  |
| Bikini Row 34 1st BLS to 2nd BLS                          | $11.6 \pm 0.4$        | $0.09 \pm 0.02$   | $5.3 \pm 0.4$     | 128.0                                 | 2.18                                  |
| Bikini Row 24 1st BLN to 2nd BLN                          | $7.8 \pm 0.2$         | $0.20 \pm 0.03$   | $3.5 \pm 0.3$     | 39.0                                  | 2.23                                  |
| Eneman NW end of Island 500-700 mR/hr area                | $209.2 \pm 9.0$       | $97.6 \pm 4.3$    | $24.0 \pm 1.5$    | 2.14                                  | 8.57                                  |
| Eneman 500-700 mR/hr area                                 | $360.9 \pm 5.9$       | $174.3 \pm 2.8$   | $45.0 \pm 1.0$    | 2.07                                  | 8.05                                  |

a Single sample error values are one-sigma, propagated, counting errors. See Appendix 20.  
 \* H.R. Not resolved by alpha spectroscopy.  
 \*\* N.S. Not significant.



## MARSHALL ISLANDS RADIOLOGICAL FOLLOWUP\*

N.A. Greenhouse and T.F. McCraw  
 Health Physics and Safety Division, Brookhaven National Laboratory, Upton, New York 11973  
 and  
 Division of Operational Safety, U.S. Energy Research and Development Administration, Washington, D.C. 20545

Abstract

In August, 1968, President Johnson announced that the people of Bikini Atoll would be able to return to their homeland. Thereafter, similar approval was given for the return of the peoples of Enewetak. These two regions, which comprised the Pacific Nuclear Testing Areas from 1946 to 1958, will probably be repopulated by the original inhabitants and their families within the next year. As part of its continuing responsibility to insure the public health and safety in connection with the nuclear programs under its sponsorship, ERDA (formerly AEC) has contracted Brookhaven National Laboratory to establish radiological safety and environmental monitoring programs for the returning Bikini and Enewetak peoples. These programs are described in the following paper. They are designed to define the external radiation environment, assess radiation doses from internal emitters in the human food chain, make long range predictions of total doses and dose commitments to individuals and to each population group, and to suggest actions which will minimize doses via the more significant pathways.

Introduction

The U.S. nuclear testing programs of the 1940s and 1950s had significant local environmental impacts on the coral atolls of Bikini and Enewetak in the Marshall Islands. The high level close-in fallout made these atolls uninhabitable for many years. Fallout from the BRAVO event, which took place at Bikini in 1954, was inadvertently deposited on the nearby atolls of Rongelap, Rongerik and Utirik. In all, some thirteen atolls in the northern Marshalls were probably affected to a greater or lesser extent by fallout from these nuclear tests. Of these, however, the most significant long term radiological impact was on the test atolls, Bikini and Enewetak, and on Rongelap Atoll.

In 1957, Rongelap was reoccupied by its original inhabitants who had been evacuated two days after BRAVO. During the past several years, definitive plans have been made to repatriate the original inhabitants of Bikini and Enewetak Atolls, and their families. It is hoped that their return can take place soon.

In order to identify radiological problems from residual radioactivity in the environment, and to provide a data base for dose predictions applicable to the returning populace, ERDA (and its predecessor, the AEC), has sponsored many radiological surveys in the Marshall Islands. These surveys began during test operations and have been conducted periodically up to the present time. Results of the surveys have been published in numerous reports and scientific journals. References 1 through 12 are published reports of AEC/ERDA supported surveys of these atolls. References 13 through 29 are a portion of the published reports on work with collected environmental samples supported by AEC/ERDA.

Evaluation of survey results for Bikini Atoll, the consideration of predicted exposures compared with applicable radiation standards, and the acknowledgement of the many benefits to the people if they could return, led to the decision to clean up and rehabilitate that atoll. The Department of Defense, Department of the Interior (DOI), and AEC (now ERDA) participated in a joint effort of clean up and rehabilitation of Bikini Atoll starting in February, 1969. Clean up was completed in the fall of that year. Agricultural rehabilitation and housing construction is being conducted by DOI.

The decision to return the Enewetakese to their atoll led to a comprehensive survey conducted at Enewetak in 1972-1973.<sup>(10)</sup> A regional survey planned for 1976 will provide baseline radiological data for future dose assessments throughout nearly all of the northern Marshall Islands which may have been affected by the testing program. Environmental evaluations at Rongelap and Utirik Atolls have been undertaken periodically in association with ERDA's medical evaluations program there over the past 20 years.<sup>(30-42)</sup>

From all of these earlier surveys, it became apparent that periodic environmental monitoring and dose assessments must be made for Bikini, Enewetak, Rongelap and perhaps other atolls in the northern Marshalls to maintain a current radiological data base and to provide current information on individual and population doses. This followup monitoring is being performed by Brookhaven National Laboratory at the request of the Division of Operational Safety, U.S. Energy Research and Development Administration.

---

\*Research carried out at Brookhaven National Laboratory under the auspices of the U. S. Energy Research and Development Administration. By acceptance of this article, the publisher and/or recipient acknowledges the U.S. Government's right to retain a nonexclusive, royalty-free license in and to any copyright covering this paper.

### Radiological Concerns

The primary radiological problems are the result of residual fission and activation products in the terrestrial environment. They have been identified by previous environmental surveys as follows: 1) External radiation levels significantly higher on some islands in an atoll compared to levels on lightly contaminated islands. 2) Fission and activation product radioactivity in certain terrestrial food items now growing on islands of these atolls and the possibility that unacceptable levels of these radionuclides may appear in foods, plants and animals newly introduced into these atolls. 3) Radioactivity in the ground water, a possible source of drinking water and water for irrigation. 4) Plutonium and americium isotopes in the surface soil. These factors are illustrated by data in Tables 1 through 4 taken from previous radiological survey reports.

| Table 1. Gamma Radiation Rates in Bikini Atoll <sup>1</sup> |                        |   |
|---|------------------------|---|
| (mR/hr)   |                        |   |
| Island  | Exposure rate<br>Range | Major contributors                                      |
| Bikini  | .010-.120              | <sup>137</sup> Cs                                       |
| Weathered areas   | .010-.030              |   |
| Close to shore  | .020-.040              |   |
| Island center   | .050-.080              |   |
| Hot spots   | .080-.120+             |   |
| Eneu  | .002-.010              | <sup>137</sup> Cs                                       |
| Nam   | .010-.330              | <sup>60</sup> Co, <sup>137</sup> Cs                     |
| Outer edge  | .010-.030              |   |
| Island center   | .015-.150              |   |
| N.E. corner   | .110-.330              |   |
| Bokantuak, Iomelan,<br>Rojkere, Eonjebi                     | .003-.010              | **  |
| Aerokoj-Emenan complex:                                     |                        |   |
| Aerokoj, Aerokojlol   | .001-.010              | **  |
| Bikdrin, Lele   | .006-.010              | **  |
| Eneman  | .001-.570              | <sup>60</sup> Co, <sup>125</sup> Sb, <sup>102m</sup> Rh |
| East Eneman   | .001-.010              |   |
| West Eneman   | .020-.570              |   |
| Enidrik   | .003-.235              | <sup>60</sup> Co, <sup>125</sup> Sb, <sup>102m</sup> Rh |
| East Enidrik  | .003-.030              |   |
| West Enidrik  | .010-.235              |   |
| Lukoj   | .060-.200              | <sup>60</sup> Co, <sup>125</sup> Sb, <sup>102m</sup> Rh |
| Jelete  | .060-.130              | **  |
| Oroken  | .015-.045              | **  |
| Bokaetoktok   | .010-.035              | **  |
| Bokdrolul   | .020-.050              |   |
| Bokbata   | .010-.030              | <sup>60</sup> Co, <sup>137</sup> Cs                     |
| Aomen-Iroi complex:   |                        |   |
| Aomen   | .005-.020              | **  |
| Lomilik   | .020-.330              | <sup>60</sup> Co, <sup>125</sup> Sb                     |
| Odrin, Iroi   | .010-.040              | **  |

\* See ref. 9.

\*\* No soil sample or field spectra measurements.

In some cases, the predicted doses and dose commitments derived from survey information for Bikini and Enewetak Atolls approach or even exceed national and international radiation protection standards for certain living and dietary patterns. Corrective actions or restrictions must be placed on use of these atolls and their resources to assure that the applicable radiation standards are not exceeded. Herein lies the primary justification for the continuing environmental followup surveys sponsored by ERDA.

### Environmental Monitoring

The most important sources of exposure to people living on Rongelap and to future residents of Bikini and Enewetak Atolls are from internal deposition of radioisotopes from certain elements in the human diet, and from the long term occupancy of islands having external radiation dose rates higher than natural background. Aside from periodic re-evaluations to establish trends in external dose rate reduction, external radiation monitoring will assume less significance, compared to monitoring of the food chain, as time passes. At present, annual visits are being made to identify and collect representative samples of local diets for laboratory analysis and dose commitment updates. New locally grown food items are becoming available in small quantities on Bikini Island as a result of the experimental agricultural practices of a small group of caretaker families living there. Neither Bikini Atoll, where radiological cleanup has been completed, nor Enewetak Atoll where clean up has not yet begun, have a subsistence agriculture resource in being which is sufficient to support the anticipated populations which will one day live there (though such crops are currently being developed or planned).

Table 2. Concentrations of  $^{40}\text{K}$ ,  $^{60}\text{Co}$ ,  $^{90}\text{Sr}$ , and  $^{137}\text{Cs}$  in Food Plants Collected on Bikini Island in November 1971 and March 1972\*

| Plant type | Tissue       | Collection date | pCi/g, dry <sup>†</sup> |                  |                   |                  |
|------------|--------------|-----------------|-------------------------|------------------|-------------------|------------------|
|            |              |                 | $^{40}\text{K}$         | $^{60}\text{Co}$ | $^{137}\text{Cs}$ | $^{90}\text{Sr}$ |
| Lettuce    | Entire       | March 1972      | 7.1±6.1                 | NS <sup>‡</sup>  | 1860±10           | 320 ±25          |
| Papaya #1  | Seeds        | March 1972      | 17 ±2                   | 0.13±0.12        | 140± 1            | NA <sup>§</sup>  |
| "          | Fruit        | March 1972      | 31 ±3                   | NS               | 160± 1            | 72 ± 3           |
| Papaya #2  | Seeds        | March 1972      | 14 ±4                   | NS               | 270± 2            | NA               |
| "          | Fruit        | March 1972      | 23 ±3                   | NS               | 290± 1            | 69 ± 4           |
| Pandanus   | Fruit        | November 1971   | 7.5±0.6                 | 0.05±0.03        | 82± 0.4           | 56 ± 0.6         |
| "          | (edible)     |                 |                         |                  |                   |                  |
| "          | Fruit        | November 1971   | 12 ±2                   | NS               | 200± 1            | 220 ± 2.2        |
| "          | (fibrous)    |                 |                         |                  |                   |                  |
| "          | Leaves       | November 1971   | 3.4±1.5                 | NS               | 71± 0.4           | 190 ± 1.9        |
| Coconut #1 | Meat         | November 1971   | 1.6±1.3                 | NS               | 93± 0.5           | NA               |
| "          | Milk         | November 1971   | 3.9±1.9                 | NS               | 110± 0.7          | NA               |
| Coconut #2 | Meat         | November 1971   | 4.4±2.1                 | NS               | 110± 1            | 0.88± 0.04       |
| "          | Milk         | November 1971   | 4.3±2.5                 | NS               | 100± 1            | < .22            |
| Coconut #3 | Meat         | November 1971   | 11 ±4                   | NS               | 147± 1            | NA               |
| Coconut #4 | Meat         | November 1971   | 2.5±2.0                 | NS               | 100± 0.8          | NA               |
| "          | Milk         | November 1971   | 3.0±1.8                 | NS               | 77± 0.6           | NA               |
| Coconut #5 | Meat         | November 1971   | 15 ±6                   | NS               | 270± 2            | NA               |
| "          | Milk         | November 1971   | 2.1±1.3                 | NS               | 33± 0.3           | NA               |
| Coconut    | Fronds (old) | November 1971   | 7.0±5.0                 | NS               | 310± 3            | NA               |
|            | Fronds (new) | November 1971   | 14 ±5                   | NS               | 220± 2            | NA               |

\* See ref. 11.

<sup>†</sup>The error terms for  $^{40}\text{K}$ ,  $^{60}\text{Co}$ , and  $^{137}\text{Cs}$  are two-sigma, propagated, counting errors. The errors for  $^{90}\text{Sr}$  are one-sigma, propagated, counting errors.

<sup>‡</sup>NS = not significant. The net sample count is less than the two-sigma, propagated, counting error.

<sup>§</sup>NA = not analyzed.

Table 3. Some Radionuclides in Water Samples Collected with a Large Volume Filter Sorption Bed from Bikini Atoll, May 1972\*

| Collection location                            | Fraction                 | Liters Sampled | Radionuclide concentration in pCi/m <sup>3</sup> <sup>†</sup> |                   |                   |                   |                   |
|--|--------------------------|----------------|---|-------------------|-------------------|-------------------|-------------------|
|  |                          |                | $^{60}\text{Co}$  | $^{137}\text{Cs}$ | $^{155}\text{Eu}$ | $^{207}\text{Bi}$ | $^{241}\text{Am}$ |
| Bravo Crater (bottom)                          | Particulate <sup>‡</sup> | 3785           | 51 ±3   | 12 ±1             | 97 ±4             | 27 ± 2            | 70 ±5             |
|  | Soluble <sup>§</sup>     | 3785           | 28 ±8   | <14 ±             | <20               | 160 ±11           | <30               |
| Bravo Crater (surface)                         | Particulate              | 3785           | 6.5±1.4   | NS <sup>‡</sup>   | 9.6±1.3           | 0.5± 0.3          | 8.2±1.4           |
|  | Soluble                  | 3785           | <10   | < 6               | <16               | <12               | <22               |
| Bokdrolul Pass (ebb tide)                      | Particulate              | 4088           | 6.0±7.3   | NS                | 6.4±1.2           | NS                | 5.4±1.1           |
|  | Soluble                  | 4088           | 1.4±0.6   | <1.0              | <2.3              | 3.0± 0.5          | <2.9              |
| Bokdrolul Pass (flood tide)                    | Particulate              | 4921           | 2.1±0.7   | NS                | 1.5±0.8           | NS                | 1.8±0.9           |
|  | Soluble                  | 4921           | NS  | NS                | NS                | 2.5± 1.7          | NS                |
| Bikini Island (seaward reef)                   | Particulate              | 1620           | 1.5±1.5   | NS                | NS                | 2.0± 0.8          | NS                |
|  | Soluble                  | 1620           | 6.2±5.4   | NS                | NS                | NS                | NS                |
| Bikini Island (lagoon)                         | Particulate              | 2271           | 5.6±1.0   | NS                | NS                | 0.76± 0.38        | 1.1±1.1           |
|  | Soluble                  | 2271           | 9.2±6.6   | NS                | NS                | NS                | NS                |
| Ocean between Bikini and Enewetak <sup>§</sup> | Particulate              | 4898           | NS  | 7.7±1.0           | NS                | NS                | NS                |
|  | Soluble                  | 4898           | NS  | NS                | NS                | NS                | NS                |
| Bikini Island (freshwater well)                | Particulate              | 1893           | 21 ±3   | 21 ±1             | 14 ±3.7           | NS                | NS                |
|  | Soluble                  | 1893           | 54 ±9   | 990 ±60           | <2                | <7                | 34 ±0             |

\* See ref. 11.

<sup>†</sup>Errors are two-sigma, propagated, counting errors.

<sup>‡</sup>Particulate--that portion retained by the 0.3  $\mu$  filter.

<sup>§</sup>Soluble--that portion which passes through the 0.3  $\mu$  filter and is sorbed by the  $\text{Al}_2\text{O}_3$  beds.

<sup>‡</sup>NS--not significant. The net sample count is less than the two-sigma, propagated counting error.

<sup>§</sup>This sample was collected over a 6 hr period between the following positions: 11°29'5" by 164°58'0" E to 11°24'5" N by 164°18'0" E.

Table 4.  $^{239,240}\text{Pu}$ ,  $^{238}\text{Pu}$  and  $^{241}\text{Am}$  in Surface Soil Samples Collected at Bikini Atoll in 1972, activities in  $\text{pCi/g} \pm 1\sigma$ \*

| Location   | $^{239,240}\text{Pu}$ | $^{238}\text{Pu}$ | $^{241}\text{Am}$ | $^{239,240}\text{Pu}/^{238}\text{Pu}$ | $^{239,240}\text{Pu}/^{241}\text{Am}$ |
|--|-----------------------|-------------------|-------------------|---------------------------------------|---------------------------------------|
| Isl-29, Ourukaen   | 16.3 $\pm$ 0.5        | 5.7 $\pm$ 0.2     | 3.6 $\pm$ 0.3     | 2.86                                  | 4.53                                  |
| Isl-30, Bokoatokutoku  | 15.1 $\pm$ 0.7        | 3.9 $\pm$ 0.3     | 4.3 $\pm$ 0.42    | 3.87                                  | 3.51                                  |
| Isl-30, Bokoatokutoku<br>Pisonia Grove                             | 22.2 $\pm$ 0.5        | 6.7 $\pm$ 0.2     | 7.0 $\pm$ 0.30    | 3.31                                  | 3.17                                  |
| Boro Bokororyruru, Isl-31<br>down center of island                 | 36.4 $\pm$ 2.8        | 7.2 $\pm$ 0.8     | 13.0 $\pm$ 1.1    | 5.05                                  | 2.75                                  |
| Namu, west end + 150 yds   | 24.0 $\pm$ 0.1        | 0.28 $\pm$ 0.02   | 14.0 $\pm$ 0.4    | 85.7                                  | 1.68                                  |
| Namu, 200 yds SW of bunker<br>300 yds E of west tip                | 20.1 $\pm$ 0.3        | 0.24 $\pm$ 0.02   | 11.0 $\pm$ 0.7    | 83.8                                  | 1.76                                  |
| Namu, top of bunker center<br>of island                            | 22.9 $\pm$ 0.7        | 0.31 $\pm$ 0.04   | 15.0 $\pm$ 0.05   | 73.9                                  | 1.57                                  |
| Namu, 200 yds E of bunker,<br>center of island                     | 17.4 $\pm$ 0.6        | 0.57 $\pm$ 0.11   | 10.0 $\pm$ 0.5    | 30.5                                  | 1.68                                  |
| Bikini, Row 24 center BL to<br>1st BLN                             | 3.3 $\pm$ 0.1         | 0.45 $\pm$ 0.04   | 2.2 $\pm$ 0.3     | 7.33                                  | 1.50                                  |
| Bikini, N corner of ctr. BL<br>and Lagoon Beach Rd.                | 3.41 $\pm$ 0.36       | N.R. <sup>†</sup> | .87 $\pm$ 0.12    | --                                    | 4.01                                  |
| Bikini, Row 34 center BL to<br>1st BL                              | 3.0 $\pm$ 0.2         | 0.06 $\pm$ 0.04   | 2.1 $\pm$ 0.2     | 50.0                                  | 1.42                                  |
| Bikini, Row 38 2nd BLN to<br>Lagoon Beach Rd.                      | 2.5 $\pm$ 0.2         | 0.07 $\pm$ 0.04   | 1.2 $\pm$ 0.2     | 35.7                                  | 2.08                                  |
| Bikini, Row 25 or 26 sand-<br>pile sample, 100 yds S<br>of 2nd BLN | 0.50 $\pm$ 0.05       | N.S. <sup>‡</sup> | --                | --                                    | --                                    |
| Bikini, Row 34 ctr BL to<br>1st BLS                                | 10.8 $\pm$ 0.04       | N.R.              | 3.3 $\pm$ 0.3     | --                                    | 3.27                                  |
| Bikini, Row 24 ctr BL to<br>1st BLS                                | 13.2 $\pm$ 0.3        | N.R.              | 8.4 $\pm$ 0.55    | --                                    | 1.58                                  |
| Bikini, Row 24, 1st BLN to<br>Lagoon Beach Rd.                     | 9.3 $\pm$ 0.4         | 0.39 $\pm$ 0.07   | 4.1 $\pm$ 0.2     | 23.8                                  | 2.27                                  |
| Bikini, Row 34, 1st BLS to<br>2nd BLS                              | 11.6 $\pm$ 0.4        | 0.09 $\pm$ 0.02   | 5.3 $\pm$ 0.4     | 128.0                                 | 2.18                                  |
| Bikini, Row 24, 1st BLN to<br>2nd BLN                              | 7.8 $\pm$ 0.2         | 0.20 $\pm$ 0.03   | 3.5 $\pm$ 0.3     | 39.0                                  | 2.23                                  |
| Eneman, NW end of island<br>500-700 mR/hr area                     | 209.2 $\pm$ 9.0       | 97.6 $\pm$ 4.3    | 24.0 $\pm$ 1.5    | 2.14                                  | 8.57                                  |
| Eneman, 500-700 mR/hr area   | 360.9 $\pm$ 5.9       | 174.3 $\pm$ 2.8   | 45.0 $\pm$ 1.0    | 2.07                                  | 8.05                                  |

\* Single sample error values are one-sigma, propagated, counting errors. See ref. 29.

† N.R. Not resolved by alpha spectroscopy.

‡ N.S. Not significant.

As a result, some of the dietary items likely to have the higher radionuclides content, e.g. pandanus and breadfruit, are not actual problems to date. They may or may not be of concern in the future as the plantings mature and the fruit becomes available in quantity. Thus, the diets of people living in these two atolls are expected to change over the coming years reflecting the relative influences of imported and locally grown food items. Allowance has been made for this in development of radiation dose estimates. Experimental studies at Enewetak may yield techniques to interrupt or break the recycling of radionuclides through the vegetation, soil, and ground water systems, and thereby reduce the radioactivity content of some important dietary items. All of the aforementioned factors will necessitate continuing monitoring of the diet for many years. Periodic sampling and analysis of soil and ground water will be necessary in order to establish trends in the changes of radioactivity content of these media.

In the northern Marshalls, drinking water is obtained primarily from rain water catchments. While the radionuclide content of collected rain water will not be zero, this source is not expected to contribute significantly to the radiation exposure picture for future Bikini, Enewetak, and Rongelap Atoll residents. However, rain water which drains from the windward side of building rooftops may provide useful data on resuspension of radioactivity in the soil. The collection of rain water by future Bikini and Enewetak residents is being facilitated by including gutters and water storage tanks in plans for houses and community structures. Some of the larger islands have fresh ground water located only a few feet below the surface. Analysis of this water for its radionuclide content has been limited to date and the capacity of this resource to serve the needs of island residents is not well defined. More study of this water is being supported by ERDA.

#### Personnel Monitoring

Dose predictions for Bikini and Enewetak Atoll residents derived from environmental data have been deliberately conservative, and establish probable upper limits on doses to be expected for individuals.

Reliable assessments of actual doses must be determined through personnel monitoring. External radiation dosimeters do not appear to be a practical means of personnel monitoring for individual external dose measurements, although certain individuals within given populations may be relied upon to wear them. A "lifestyle model" which includes estimates of occupancy factors for various locations in a given atoll has been coupled with environmental monitoring data to estimate average external radiation doses to individuals. This model will be revised as needed so that it closely approximates the actual lifestyle of the people.

The more important internal pathway can be monitored directly by conventional techniques of bioassay and whole body counting of individuals. A portable shadow shield whole body counter has been constructed and mounted in a shipboard trailer for use in the Marshall Islands. It is capable of quantitative detection of very small quantities of certain radionuclides in the body such as  $^{137}\text{Cs}$  and  $^{60}\text{Co}$ , the primary environmental gamma emitters at Bikini, Enewetak and Rongelap Atolls. The system clearly identifies individuals in the Rongelap population who are not following the recommended dietary restrictions on eating coconut crabs from certain locations.<sup>(42,43)</sup> Body burdens of  $^{90}\text{Sr}/^{90}\text{Y}$ ,  $^{239,240}\text{Pu}$  and  $^{241}\text{Am}$  are estimated by the radiochemical analysis of urine samples. Urine sample collections and whole body counting will be performed every one to two years at Bikini and Enewetak Atolls when the people return, and every two to three years at Rongelap Atoll until the results warrant less frequent measurement intervals.

#### Summary

Marshall Islands Radiological Followup has consisted of intensive environmental studies at Bikini, Enewetak, and Rongelap Atolls to gather radiological data on the external radiation environment and on radioactivity in food chains. Radiation and radioactivity levels in these atolls are being reduced with time. These changes are monitored in annual or biannual environmental surveys. Updated information is used to make conservative estimates of population doses and dose commitments. When people have returned, actual internal doses to individuals are determined for whole body counting and bioassay data. These results are combined with environmental data on the external radiation environment to complete the total dose assessment picture.

#### References

1. Dunning, G. M., "Radioactive Contamination of Certain Areas in the Pacific Ocean from Nuclear Tests, A Summary of the Data from the Radiological Surveys and Medical Examinations", USAEC Report, August, 1957.
2. Held, E. E., "Gamma Dose Rates at Rongelap Atoll, 1954-1963", USAEC Report, UWFL-91, May 1965.
3. Donaldson, L. R., et al. "Bikini-Eniwetok Studies, 1964, Part I and II", University of Washington, Laboratory of Radiation Biology, UWFL-93, September 15, 1966.
4. Beck, H. L., Bennett, B. G. and McCraw, T. F., "External Radiation Levels on Bikini Atoll", May 1967, USAEC Report, HASL-190, December 1967.
5. Bennett, B. G. and Beck, H. L., "External Radiation on Bikini Atoll", Nature 223: 925-928, 1969.
6. Held, E. E., "Radiological Resurvey of Animals, Soils and Ground Water at Bikini Atoll, 1969-1970", University of Washington, College of Fisheries, NVO-269-8 (Rev. 1), February 1971.
7. Smith, A. E. and Moore, W. E., "Report of Radiological Clean up of Bikini Atoll, Office of Dose Assessment and Systems Analysis", Western Environmental Research Laboratory, Environmental Protection Agency, SWRHL-111r, January 1972.
8. McCraw, T. F. and Lynch, O. D. T. Jr., "Exposure Rate Reduction on Bikini Island Due to Concrete Dwellings, USAEC Report, WASH-1273, June 1973.
9. McCraw, T. F., "Levels of Environmental Radioactivity in Bikini Atoll", USAEC Report, WASH-1289, 1974.
10. Enewetak Radiological Survey, Volumes I, II and III, USAEC Nevada Operations Office, Report NVO-140, October 1973.
11. Lynch, O. D. T. Jr., McCraw, T. F., Nelson, V. A., and Moore, W. E., "Radiological Resurvey of Food, Soil, Air and Ground Water at Bikini Atoll, 1972", USERDA Report, ERDA-34, February 1975.
12. Gudiksen, P. H., and Robison, W. L., "Preliminary External Dose Estimates for Future Bikini Atoll Inhabitants, Lawrence Livermore Laboratory Report, Preliminary UCRL-51879, August 1975.
13. Palumbo, R. F. and Lowman, F. G., "The Occurrence of Antimony-125, Europium-155, Iron-55 and Other Radionuclides in Rongelap Atoll Soil", USAEC Report UWFL-56, April 7, 1958.
14. Lowman, F. G., "Marine Biological Investigations at the Eniwetok Test Site", from Disposal of Radioactive Wastes, International Atomic Energy Agency, Vienna, 1960.
15. Walker, R. B., Held, E. E. and Gessel, S. P., "Radiocesium in Plants Grown on Rongelap Atoll Soils, Rec. Advan. Bot. 2: 1363-1367, 1961.
16. Cole, D. W., Gessel, S. P. and Held, E. E., "Tension Lysimeter Studies of Ion and Moisture Movement in Glacial Till and Coral Atoll Soils", Soil Sci. Soc. Am. Proc. 25: 321-325, 1961.
17. Lowman, F. G. and Palumbo, R. F., "Occurrence of Bismuth-207 at Eniwetok Atoll", Nature 193: 796-797, 1962.
18. Chakravarti, D. and Held, E. E., "Chemical and Radiochemical Composition of the Rongelapese Diet", J. Food Sci. 28: 221-228, 1963.
19. Kenady, R. M., "The Soils of Rongelap Atoll, Marshall Islands", University of Washington Press, Corvallis, 1962.
20. Held, E. E., "Qualitative Distribution of Radionuclides at Rongelap Atoll", Amer. Inst. Biol. Sci., Reinhold Publ. Co., New York, 1963, pp. 167-169.

21. Held, E. E., Gessel, S. P., Marson, L. J. and Billings, R. F., "Autoradiography of Sectioned Soil Cores, Radioisotope Sample Measurement Techniques in Medicine and Biology"; International Atomic Energy Agency, Vienna, 1965.
22. Held, E. E., Gessel, S. P. and Walker, R. B., "Atoll Soil Types in Relation to the Distribution of Fallout Radionuclides", USAEC Report, UWFL-92, August 1965.
23. Welander, S. D., "Distribution of Radionuclides in the Environment of Eniwetok and Bikini Atolls, August 1964", in Proc. 2nd Natl. Symp., Radioecology, CONF-670503, Oak Ridge, Tenn., 1967.
24. Beasley, T. M. and Held, E. E., "Nickel-63 in Marine and Terrestrial Biota, Soil, and Sediment", Science 164: 1161-1163, 1969.
25. Beasley, T. M. and Held, E. E., "Silver-108m in Biota and Sediments at Bikini and Eniwetok Atolls", Nature 230: 450-451, 1971.
26. Schell, W. R. and Yang, A. I. C., "Long-Lived Radionuclides Produced at Bikini and Eniwetok Atolls"; University of Washington, College of Fisheries, RLO-2225-T18-3, April 1973.
27. Schell, W. R., "Studies of Concentrations of Unreported Long-Lived Radionuclides in Biota and Ocean Sediments at Bikini and Eniwetok Atolls"; University of Washington, College of Fisheries, RLO-2225-T18-4, May 1973.
28. Noskin, V. E., Wong, K. M., Eagle, R. J. and Gatrousis, C., Lawrence Livermore Laboratory Report, UCRL-51612, June 1974.
29. Nevissi, A., Schell, W. R. and Nelson, V. A., "Plutonium and Americium in Soils of Bikini Atoll" in IAEA/ERDA Symp. on Transuranium Nuclides in the Environment, San Francisco, RLO-2225-T18-17, in press.
30. Cronkite, E. P., et al., "Some Effect of Ionizing Radiation in Human Beings: A Report on the Marshallese and American Accidentally Exposed to Radiation Fallout and a Discussion of Radiation Injury in the Human Being", AEC-TID 5385, U.S. Govt. Printing Office, Washington, D. C., 1956.
31. Bond, V. P., Conard, R. A., Robertson, J. S. and Weden, E. A. Jr., "Medical Examination of Rongelap People Six Months after Exposure to Fallout", WT-937, Operation Castle Addendum Report 4.1A, April 1955.
32. Cronkite, E. P., et al., "Twelve-Month Postexposure Survey on Marshallese Exposed to Fallout Radiation, BNL 384 (T-71), 1955.
33. Conard, R. A., et al., "Medical Survey of Marshallese Two Years after Exposure to Fallout Radiation, J. Amer. Med. Assoc. 164: 1192, 1957.
34. Conard, R. A., et al., "March 1957 Medical Survey of Rongelap and Utirik People Three Years After Exposure to Radioactive Fallout"; BNL 501 (T-119), June 1958.
35. Conard, R. A., et al., "Medical Survey of Rongelap People, March 1958, Four Years After Exposure to Fallout, BNL 534 (T-135), May 1959.
36. Conard, R. A., et al., "Medical Survey of Rongelap People Five and Six Years After Exposure to Fallout, BNL 609 (T-179), September 1960.
37. Conard, R. A., et al., "Medical Survey of Rongelap People Seven Years After Exposure to Fallout", BNL 727 (T-260), May 1962.
38. Conard, R. A., et al., "Medical Survey of Rongelap People Eight Years After Exposure to Fallout", BNL 780 (T-296), January 1963.
39. Conard, R. A., et al., "Medical Survey of the People of Rongelap and Utirik Islands Nine and Ten Years After Exposure to Fallout Radiation (March 1963 and March 1964), BNL 908 (T-371), May 1965.
40. Conard, R. A., et al., "Medical Survey of the People of Rongelap and Utirik Islands Eleven and Twelve Years After Exposure to Fallout Radiation (March 1965 and March 1966), BNL 50029 (T-446) April 1967.
41. Conard, R. A., et al., "Medical Survey of the People of Rongelap and Utirik Islands Thirteen, Fourteen, and Fifteen Years After Exposure to Fallout Radiation (March 1967, March 1968 and March 1969), BNL 50220 (T-562), June 1970.
42. Conard, R. A., et al., "A Twenty-Year Review of Medical Findings in a Marshallese Population Accidentally Exposed to Radioactive Fallout", BNL50424, September 1975.
43. Cohn, S. H., Conard, R. A., Gusmano, E. A., and Robertson, J. S., "Use of a Portable Whole Body Counter to Measure Internal Contamination in a Fallout-Exposed Population, Health Phys. 9: 15 (1963).